



Review article

Promoting brain health through physical activity among adults exposed to early life adversity: Potential mechanisms and theoretical framework

Shannon D. Donofry^{a,b,*}, Chelsea M. Stillman^a, Jamie L. Hanson^{a,c,d}, Margaret Sheridan^e, Shufang Sun^{f,g}, Eric B. Loucks^{f,g,h}, Kirk I. Erickson^{a,d,i,j,k}

^a Department of Psychology, University of Pittsburgh, Pittsburgh, PA, United States

^b Psychiatric and Behavioral Health Institute, Allegheny Health Network Pittsburgh, PA, United States

^c Learning Research and Development Center, University of Pittsburgh, Pittsburgh, PA, United States

^d Center for the Neural Basis of Cognition, Pittsburgh, PA, United States

^e Department of Psychology and Neuroscience, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States

^f Department of Behavioral and Social Sciences, Brown University School of Public Health, Providence, RI, United States

^g Mindfulness Center, Brown University, Providence, RI, United States

^h Department of Epidemiology, Brown University School of Public Health, Providence, RI, United States

ⁱ Center for Neuroscience, University of Pittsburgh, Pittsburgh, PA, United States

^j Murdoch University, College of Science, Health, Engineering, and Education, Perth, Western Australia, Australia

^k PROFITH "PROMoting FITness and Health Through Physical Activity" Research Group, Department of Physical Education and Sports, Faculty of Sport Sciences, University of Granada, Granada, Spain



ARTICLE INFO

Keywords:

Early life adversity
Adverse childhood experiences
Physical activity
Brain health
Midlife
Aging
Reversibility network

ABSTRACT

Adverse childhood experiences such as abuse, neglect, and poverty, profoundly alter neurobehavioral development in a manner that negatively impacts health across the lifespan. Adults who have been exposed to such adversities exhibit premature and more severe age-related declines in brain health. Unfortunately, it remains unclear whether the negative effects of early life adversity (ELA) on brain health can be remediated through intervention in adulthood. Physical activity may represent a low-cost behavioral approach to address the long-term consequences of ELA on brain health. However, there has been limited research examining the impact of physical activity on brain health among adults with a history of ELA. Accordingly, the purpose of this review is to (1) review the influence of ELA on brain health in adulthood and (2) highlight evidence for the role of neurotrophic factors, hypothalamic-adrenal-pituitary axis regulation, inflammatory processes, and epigenetic modifications in mediating the effects of both ELA and physical activity on brain health outcomes in adulthood. We then propose a theoretical framework to guide future research in this area.

1. Introduction

Early life adversity (ELA) encompasses a range of negative experiences during childhood and adolescence, including poverty, abuse, and parental maladjustment which interfere with neurocognitive and socioemotional development. Unfortunately, ELA is common throughout the world, prevalent in both economically distressed and wealthy countries. It is estimated that 40–60 % of individuals are exposed to at least one form of ELA, and a quarter of individuals experience multiple adversities (Felitti et al., 1998; Kessler et al., 2010; Merrick et al., 2018). Notably, marginalized and underserved populations such as racial and

ethnic minorities, sexual and gender minorities, and low-income communities are overburdened with early life adversity (Merrick et al., 2018), contributing to health inequities in adulthood.

Exposure to ELA exerts profound and long-lasting effects on health and well-being that persist into late adulthood. Globally, ELA accounts for up to 30 % of incident psychiatric disorders in adults (Kessler et al., 2010), and increases risk for the development of numerous chronic health conditions, such as Type II diabetes, obesity, and cardiovascular disease (Friedman et al., 2015; Merrick et al., 2019; Su et al., 2015). More recent evidence suggests that ELA also negatively impacts brain health in mid- and late life, accelerating the rate of age-related cognitive

* Corresponding author at: Department of Psychology, University of Pittsburgh, Clinical Health Psychologist, Psychiatry and Behavioral Health Institute, Allegheny Health Network, 4 Allegheny Center, 8th Floor, Pittsburgh, PA, 15212, United States.

E-mail addresses: sdd14@pitt.edu, shannon.donofry@ahn.org, shannon.donofry@gmail.com (S.D. Donofry).

<https://doi.org/10.1016/j.neubiorev.2021.09.051>

Received 1 June 2021; Received in revised form 13 September 2021; Accepted 28 September 2021

Available online 5 October 2021

0149-7634/© 2021 Elsevier Ltd. All rights reserved.

decline and heightening susceptibility to neurological illnesses such as Alzheimer's Disease (AD; [Short and Baram, 2019](#)). Further, retrospective studies have observed a link between ELA and brain structure and function in adulthood ([Botros et al., 2019](#); [Pechtel and Pizzagalli, 2011](#)). These data suggest that the effects of ELA on brain health can be long-lasting and may underlie ELA-related susceptibility to psychological, cardiometabolic, and neurocognitive illness. These issues raise questions regarding whether and to what degree the negative effects of ELA on brain health in adulthood can be mitigated or reversed. In an effort to address such questions, the National Institute on Aging formed the Reversibility Network, an interdisciplinary research network focused on advancing understanding of the mechanisms of ELA-related declines in health in later life to facilitate the development of behavioral interventions to remediate or reverse these effects. The present review was guided by and written in the service of the goals of the Reversibility Network.

Physical activity may be an effective low-cost and widely accessible behavioral approach to remediate the long-term negative health consequences of exposure to ELA. In addition to the well-documented benefits of physical activity for reducing chronic disease risk ([Durstine et al., 2013](#)), there is growing evidence that physical activity promotes improvements in brain health across the lifespan ([Erickson et al., 2019](#); [Hillman et al., 2008](#)), including enhanced psychological functioning and cognitive performance as well as alterations in brain structure and function ([Erickson et al., 2019](#)). Such benefits have been observed even among adults who have already experienced a decline in brain health due to illness or aging ([Chirles et al., 2017](#); [Fong et al., 2012](#); [Lautenschlager et al., 2008](#); [Oberlin et al., 2017](#); [Smith et al., 2014](#)). Further, evidence drawn from an observational study in adolescents has shown that physical activity can buffer against the negative effects of childhood abuse on later internalizing symptoms ([Healy et al., 2021](#)). There is also evidence that many of the mechanistic pathways through which ELA negatively impacts brain health are also modulated by engagement in physical activity, which we outline in detail below. Thus, physical activity may enhance and promote brain health in similar areas and potentially through similar but opposing mechanisms to those through which ELA precipitates adverse brain health in adulthood. Accordingly, the purpose of this review is to (1) summarize what is known about the influence of ELA on brain health in adulthood and (2) identify the overlapping pathways whereby physical activity may stimulate improvements in brain health among adults who have been affected by ELA. We will then propose a theoretical framework to guide future research efforts integrating across traditionally separate disciplines such as exercise physiology, developmental psychology, neurology, and health neuroscience. Given that both ELA and physical activity exert widespread and complex effects on most systems of the body, this review is intentionally broad in scope, intended to identify if there is scientific justification for future research examining the efficacy of physical activity interventions for mitigating or reversing the deleterious effects of ELA on adult brain health.

2. Key definitions

2.1. Early life adversity

Adverse childhood experiences encompass a range of disadvantageous events to which a child must adapt, including psychosocial and material adversities such as parental maladjustment, abuse, neglect, discrimination, and poverty, as well as physical exposures to environmental toxins and malnutrition (for a detailed review of how to define ELA, please see ([McLaughlin, 2016](#)). For the purposes of the present review, we focus on psychosocial adversities and have adopted an inclusive definition of ELA that comprises any psychosocial adverse event or experience occurring during childhood that may negatively impact brain health across the lifespan. This includes experiences widely viewed as adverse such as physical or emotional abuse and neglect, as

well as other psychosocial adversities such as socioeconomic disadvantage, exposure to domestic or community violence, caregiver loss, caregiver criminality and/or incarceration, caregiver psychopathology, and bullying. Although exposures occurring during the prenatal period (e.g., maternal stress) have been linked to elevated lifetime risk for chronic health conditions and poor psychosocial functioning ([Graignic-Philippe et al., 2014](#); [Tobi et al., 2018](#)), we have limited the current review to postnatal adversities from birth to age 18. We broadly refer to these experiences as “early life adversities” (ELA), though occasionally examine the impact of specific forms of ELA when warranted (e.g., multiple lines of converging evidence to support an impact on brain health).

2.2. Brain health

Similar to our approach to operationalizing ELA, we have also conceptualized brain health broadly to include brain structure and function as well as the cognitive and psychological processes that arise from brain function (e.g., executive functions, mood). This approach is consistent with the definition adopted by the 2018 Physical Activity Guidelines Scientific Advisory Committee. This committee was appointed by the U.S. Department of Health and Human Services to compile scientific evidence supporting beneficial effects of physical activity on the brain in the service of developing public health recommendations for preserving brain health across the lifespan ([Erickson et al., 2019](#); [Piercy et al., 2018](#)). Thus, brain health as we have defined it includes normative variation in mood and cognition, as well as clinical syndromes characterized by deficits in these domains (i.e., psychiatric and neurological conditions). We also include in our definition of brain health normative and pathological variation in brain structure and function, as measured predominantly using structural and functional magnetic resonance imaging.

2.3. Physical activity

In keeping with the 2018 Physical Activity Guidelines ([Erickson et al., 2019](#)), we have defined physical activity broadly to include any bodily movement that results in energy expenditure, regardless of the form (e.g., aerobic vs. anaerobic vs. strengthening), intensity (i.e., light, moderate, or vigorous intensity), and duration (e.g., acute bursts of activity vs. habitual activity sustained over days, weeks, or months). Where indicated, we have specified the effects of these different parameters of physical activity on brain health, but otherwise refer to physical activity in general terms. Further, although there are numerous studies linking cardiorespiratory fitness with indicators of brain health ([Boots et al., 2015](#); [Kramer and Colcombe, 2018](#); [Weinstein et al., 2012](#)) and cardiorespiratory fitness is a proximal outcome of regular physical activity ([Lang et al., 2018](#)), we have generally excluded these studies unless there was concomitant measurement of physical activity.

3. Mechanistic models of early life adversity

There are numerous approaches to conceptualizing the pathways through which ELA impacts health and well-being across the lifespan, each with unique assumptions and methodologies ([McLaughlin and Sheridan, 2016](#)). Examples of prevailing theoretical frameworks include cumulative risk, dimensional, and accelerated aging models. Cumulative risk approaches to classifying ELA tally the number of ELA exposures over the course of childhood and adolescence to create a cumulative risk score ([Evans et al., 2013](#)). The cumulative risk framework has been widely used to quantify ELA and determine its neurodevelopmental consequences, emphasizing in particular the influence of ELA on the development of stress response systems. Of note, this approach does not distinguish among subtypes of adverse events, nor does it typically account for duration, severity, or developmental timing of exposure, potentially obscuring important variation in exposure and the impact of

that variability on outcomes in adulthood. More recent approaches have developed and expanded upon the cumulative risk method by defining ELA along core dimensions, each of which encompasses numerous experiences that vary in severity (McLaughlin and Sheridan, 2016). An example of a dimensional model includes the dimensional model of adversity & psychopathology (DMAP; McLaughlin and Sheridan, 2016), which distinguishes between experiences that involve threat of harm to a child and those that involve the absence of expected inputs. A second example is the harshness-unpredictability model, which characterizes the degree to which a child's environment does not meet physical and socioemotional needs (e.g., physical abuse, poverty) and/or is frequently changing in a manner that a child cannot predict (e.g., frequent housing relocation, changes in family composition; Ellis et al., 2009). Dimensional models highlight additional mechanisms linking ELA to physical and psychological health outcomes beyond stress reactivity and programming, such as learning in the context of developmental plasticity. Thus, dimensional models provide additional avenues through which to develop mechanism-informed treatments aimed at preventing the downstream consequences of adversity. Finally, accelerated aging models focus on the role of ELA in programming premature aging of multiple organ systems (e.g., pulmonary, cardiovascular, renal etc.; Belsky et al., 2015) and cellular processes (e.g., advanced deterioration of telomeres and premature increases in DNA methylation; Ridout et al., 2018; Sumner et al., 2019). According to this framework, ELA prompts accelerated decline of these systems, leading to increased risk for and earlier onset of age-related disease and mortality (Shalev and Belsky, 2016). These accelerated aging models often focus on harshness-unpredictability domains of ELA, but they are not theoretically incompatible with cumulative risk or other dimensional models. Rather, accelerated aging models shift focus from childhood and adolescent outcomes to outcomes in adulthood, with an emphasis on identifying novel mechanisms of aging that may be targeted through behavioral and pharmacological interventions delivered earlier in the lifespan. Each of these frameworks has contributed to furthering scientific understanding of the short and long term sequelae of ELA. Given that research focused on adult outcomes has emerged more recently compared to research focused on the consequences of ELA in childhood and adolescence, these conceptual models have not been as thoroughly examined in the context of adulthood, and few studies in adults specify whether the hypotheses and approach were guided by a particular framework. Nevertheless, we include them here as a means of introducing the prevailing theories regarding the mechanisms through which ELA impacts brain development and brain health and to highlight the complexities inherent to this type of research.

4. Influence of early life adversities on brain health in adulthood

Early life experiences can influence brain health across the lifespan. Healthy brain development in both animals and humans is shaped by early life experiences (Fox et al., 2010; Knudsen, 2004; McLaughlin et al., 2014; Sheridan and McLaughlin, 2014). For example, responsive and dependable caregivers and environmental predictability foster a trajectory of normal brain development. In contrast, adverse childhood experiences such as neglect, abuse, or extreme poverty can interrupt normal brain development and set an individual up for a trajectory of aberrant brain, behavioral, and emotional development across the lifespan. Given the ethical challenges with manipulating ELA in humans, nearly all experimental work focused on ELA and brain health has been conducted using animal models. These models provide convincing evidence that exposure to ELA during sensitive periods disrupt typical brain development (Danese and McEwen, 2012; Bruce S. McEwen, 2003; Short and Baram, 2019) and these findings are supported by the small number of experimental studies in humans (e.g., Bick et al., 2015; Nelson et al., 2007). Although the human literature is limited predominantly to correlational and prospective studies, mounting evidence

from these studies corroborates many of the patterns observed in experimental animal models. Below we highlight the most salient patterns that have emerged in this area and summarize these effects in Fig. 1.

4.1. Psychological outcomes

ELA has been associated with impaired psychological functioning in adulthood. Adults who have been exposed to ELA are more likely to be diagnosed with a psychiatric illness (Heinonen et al., 2018), including major depressive disorder (Merrick et al., 2019; Nanni et al., 2012), anxiety disorders (Jonker et al., 2017), bipolar disorder (Bruni et al., 2018; Shapero et al., 2017), and substance use disorders (LeTendre and Reed, 2017), among others (Bruni et al., 2018; Fuller-Thomson and Lewis, 2015; Hock et al., 2018; McLaughlin et al., 2017). There is also evidence that women exposed to ELA are at elevated risk for peripartum depression, anxiety, and substance use (Currie et al., 2020; Gonzalez et al., 2009; Heldreth et al., 2016; Jonas et al., 2013; Racine et al., 2020; Young-Wolff et al., 2018). Of note, direct and indirect experience of racism during childhood has been shown to predict more severe depressive symptoms during the postpartum period among Black and African American mothers (Heldreth et al., 2016), highlighting the importance of examining this specific form of adversity when evaluating adult health outcomes. Further, ELA is related to earlier onset and more severe course of illness (Heinonen et al., 2018), particularly among adults who experienced abuse or neglect (Carbone et al., 2019; Fuller-Thomson and Lewis, 2015). ELA also prospectively predicts higher likelihood of attempting suicide in adulthood (Wanner et al., 2012), thus contributing to psychiatric mortality. Unfortunately, a history of ELA reduces the efficacy of behavioral and psychopharmacological treatments for psychiatric illness (Nanni et al., 2012), prolonging suffering and potentially increasing susceptibility for other psychiatric and medical comorbidities. Moreover, exposure to ELA may lower the threshold of severity at which a stressful life experience in adulthood precipitates onset or recurrence of clinically elevated psychiatric symptoms (Shapero et al., 2017). ELA has also been associated with a reduced capacity to regulate negative emotions as evidenced by observations that individuals with a history of ELA tend to rely on ineffective emotion regulation strategies such as suppression and rumination (Cameron et al., 2018; Hong et al., 2018). Insufficient emotion regulation skills may therefore be a key pathway by which ELA heightens vulnerability to psychological distress and psychiatric illness in adulthood. Finally, ELA may have a lasting impact on interpersonal functioning in adulthood. Childhood physical and emotional abuse has been related to reduced emotional closeness with family members in mid- and late-adulthood, an effect that persisted when controlling for concurrent depressive symptomatology (Savla et al., 2013). Thus, the psychological consequences of ELA are pervasive, impacting numerous facets of socioemotional functioning in adulthood.

4.2. Cognitive outcomes

A growing body of evidence suggests that ELA disrupts cognitive processes that are critical for the maintenance of health and well-being across the lifespan (Short and Baram, 2019). Research has demonstrated that ELA confers substantial risk for the development of cognitive impairment and neurological illnesses such as AD in late life, many decades after exposure (for review, see (Danese and McEwen, 2012). Socioeconomic disadvantage in childhood has been one of the most widely studied forms of ELA with regard to the potential impact on cognitive outcomes in adulthood, perhaps due in part to the well-established relationship between educational attainment and cognitive decline in late life (Norton et al., 2014; Stern et al., 1994). Low socioeconomic status in childhood, as typically indexed by parental education and income, has been associated with worse memory performance among older adults, even among those characterized as having




Fig. 1. (A) Theoretical models of the impact of early life adversity on brain health in adulthood. (B) Influence of early life adversity and physical activity on adult brain health.

high income at the time of assessment (Marden et al., 2017). Further, childhood socioeconomic disadvantage may impact the trajectory of cognitive aging by accelerating the rate of age-related cognitive decline (Melrose et al., 2015). Similar patterns have been observed among adults who have experienced more severe forms of ELA such as neglect and maltreatment (Geoffroy et al., 2016).

Executive functions such as cognitive flexibility, inhibitory control, working memory, and planning are particularly affected by ELA, and have been consistently shown to be disrupted among adults with a history of ELA (Almanza-Sepulveda et al., 2018; Butler et al., 2017; Lovallo et al., 2013). ELA may also increase the likelihood and severity of executive dysfunction that may arise as a consequence of some medical conditions (Hawkins et al., 2020; Shanmugan et al., 2017, 2020), which may lead to poorer prognosis and lowered quality of life. Research on the effect of ELA on other cognitive domains is more limited, though some studies have documented lower performance on tests of visuospatial ability, episodic memory, language, and processing speed among adults who have been exposed to ELA (Beck et al., 2018; Duval et al., 2017; Greenfield and Moorman, 2019).

Of note, psychiatric illnesses that may emerge as a result of ELA such as major depressive disorder (MDD) have been shown to increase vulnerability to cognitive decline and dementias (Byers and Yaffe, 2011), indicating that the effect of ELA on cognitive health in adulthood is complex. Importantly, animal models examining the impact of ELA on cognitive functioning across the lifespan have yielded similar patterns (for a review, see Lesuis et al., 2018), lending support for a causal link between ELA exposure and subsequent cognitive impairment.

4.3. Brain structure and function

The majority of studies focusing on ELA-related disruption of brain circuitry have focused on neurodevelopment in childhood and adolescence, with there being comparatively fewer studies in adults. Nevertheless, just as exposure to ELA has enduring effects on cognitive and psychological functioning, emerging research has documented brain abnormalities among adults with a history of ELA. The hippocampus appears to be particularly susceptible to the effects of ELA. For instance, exposure to adversity in childhood has been associated with reduced hippocampal volume in adulthood (Dannlowski et al., 2012; Frodl et al., 2012; Gorka et al., 2014; Teicher et al., 2012; though see Khoury, Pechtel, Andersen, Teicher, & Lyons-Ruth, 2019 & Mikolas et al., 2019 for reports of opposing patterns), with this effect being most pronounced

in the glucocorticoid-sensitive CA3 and dentate gyrus subfields (e.g., Frodl et al., 2012; Teicher et al., 2012). Further, smaller hippocampal volumes prospectively predicted heightened symptoms of anxiety following the experience of a stressful life event among adults with a history of childhood maltreatment (Gorka et al., 2014), suggesting that differences in hippocampal structure may be one pathway through which ELA exposure increases psychological vulnerability in adulthood.

ELA exposure has also been linked to volume reductions in other prefrontal and subcortical regions that support cognitive and psychological processes known to be disrupted by ELA, including the orbitofrontal cortex (OFC), medial prefrontal cortex (PFC), anterior cingulate cortex (ACC), insula, amygdala, and caudate nucleus (Brooks et al., 2016; Dannlowski et al., 2012; Gorka et al., 2014; Holz et al., 2015; Underwood et al., 2019), with these effects being detectable many decades after the experience of ELA (Gheorghie et al., 2020). White matter integrity has also been shown to be disrupted among adults exposed to ELA. For example, young adults with a parent diagnosed with a substance use disorder during their childhood were shown to exhibit reduced white matter integrity in frontocortical, frontostriatal, and frontoparietal tracts (Acheson et al., 2014). Another study demonstrated that white matter integrity of the uncinate fasciculus, a major tract connecting the amygdala to the ventromedial PFC, was lower in young adults exposed to maltreatment in childhood compared to those without maltreatment histories (Hanson et al., 2015). Further, lower white matter integrity at baseline predicted elevated internalizing symptoms following subsequent exposure to a stressful life event. Unfortunately, changes in brain structure that arise in response to more severe forms of ELA such as deprivation and institutionalization may persist even when life circumstances in adulthood are otherwise salubrious (Mackes et al., 2020). This indicates that some ELA-related decrements in brain health may be more difficult to remediate and the efficacy of interventions applied in adulthood may be moderated by the severity of ELA, a hypothesis that warrants further testing.

In addition, early life adversity is associated with alterations in the function of brain regions embedded in functional networks that orchestrate cognitive and psychological processes impacted by ELA (for a detailed review please see Herzberg and Gunnar, 2020). Several studies have demonstrated that ELA is associated with reduced resting state connectivity within the default mode network (DMN) (Elton et al., 2014; Sripada et al., 2014; L. Wang et al., 2014; Zeev-Wolf et al., 2019). The DMN is a robust functional network comprised of regions in medial PFC (mPFC), temporal (e.g., hippocampus), and posterior cingulate

cortices that supports internally-guided and self-referential cognitive processes (Buckner and DiNicola, 2019; Greicius et al., 2003), and has been shown to be disrupted in many conditions linked to ELA, including MDD (Greicius et al., 2007), substance use disorder (R. Zhang and Volkow, 2019), obesity (Donofry et al., 2020a, 2020b; Tregellas et al., 2011), and AD (Greicius et al., 2004; Koch et al., 2012). Exposure to childhood trauma predicts reduced connectivity between the hippocampus, dorsomedial and dorsolateral PFC in veterans with symptoms of combat-related post-traumatic stress (Birn et al., 2017). Further, among adults with a documented history of childhood poverty, lower resting connectivity between the hippocampus and posterior cingulate cortex was related to higher basal cortisol levels (Sripada et al., 2014). Several DMN hubs, including the mPFC and hippocampus, densely express glucocorticoid receptors (McEwen et al., 1986), and may therefore be more robustly dysregulated by exposure to ELA, particularly threat-related adversities. These findings suggest that alterations within the DMN may be one pathway through which ELA influences health and well-being in mid- and late life.

Altered patterns of task-evoked activation have also been noted among adults with a history of ELA. For instance, among adults diagnosed with MDD or an anxiety disorder, those with a history of ELA demonstrated exaggerated activation in corticolimbic regions such as the ACC, superior frontal gyrus, and parahippocampal gyrus in response to facial expressions of fear and anger (Herringa et al., 2013; Peters et al., 2019). Interestingly, these patterns of activation mediated the relationship between childhood adversity and tendency to engage in rumination (Peters et al., 2019), a perseverative self-focused cognitive style that impedes the use of effective problem-solving skills (Bernstein et al., 2017). Further, one study found that the relationship between childhood socioeconomic disadvantage and amygdala reactivity to threatening facial expressions persisted even when controlling for other contemporaneous factors known to influence amygdala functioning, including adult socioeconomic status, symptoms of depression and anxiety, as well as dispositional emotionality (Gianaros et al., 2008). Similarly, ELA exposure has been linked to increased reward-related activation in the ventral striatum, putamen and ACC (Boecker et al., 2014; Boecker-Schlier et al., 2016; DelDonno et al., 2019), as well as heightened functional connectivity between reward-sensitive regions (e.g., ventral striatum, mPFC) during a monetary reward task (Hanson et al., 2018). Further, ventral striatal – dorsal medial PFC connectivity evoked by reward feedback was strongest among those reporting both ELA and recent life stress, and partially accounted for the relationship between ELA and symptoms of depression and anxiety (Hanson et al., 2018), indicating that cumulative life stress may potentiate psychopathology in adulthood by modulating reward neurocircuitry. However, other studies have observed ELA-related blunting of activation in reward-sensitive regions in both childhood (Birn et al., 2017; Mullins et al., 2020) and adulthood (Gianaros et al., 2011; Hanson et al., 2016), highlighting the complexity and heterogeneity of the neurobiological effects of ELA.

In contrast to reports of increased activation and connectivity evoked by psychological stressors and reward stimuli among adults with a history of ELA, cognitive tasks (e.g., Stroop task, Go/No-Go task) evoke less activation in cognitive, emotion, and visceral control regions such as the dorsolateral PFC (dlPFC), ACC, amygdala, and hypothalamus in ELA-exposed individuals (Banihashemi et al., 2015; Harms et al., 2017). Notably, reduced activation of these regions was associated with slowed task response times (Harms et al., 2017) and more substantial task-induced increases in heart rate (Banihashemi et al., 2015). These findings suggest that ELA may contribute to less successful recruitment of prefrontal and subcortical regions in the face of increased cognitive demands, which may contribute to ELA-related susceptibility to poor psychological, cognitive and cardiovascular outcomes.

It is important to note that many of the psychiatric and physical health conditions that have been linked with exposure to ELA have also been shown to exert deleterious effects on the structure and function of

the brain, making it difficult to disentangle the extent to which the relationship between ELA and brain outcomes are mediated by, independent of, or amplified by the effects of these conditions on the brain. Additional research is needed to clarify how ELA interacts with psychiatric and physical health conditions in adulthood to affect brain aging trajectories.

5. Influence of physical activity on brain health in adulthood

Physical activity improves many of the aforementioned negative outcomes known to be precipitated by ELA. This suggests that physical activity interventions may represent a promising approach for mitigating or reversing the negative consequences of ELA on brain health. In the following sections, we summarize the effects of physical activity on brain health with a particular focus on how physical activity may influence outcomes associated with ELA.

5.1. Psychological outcomes

Evidence from prospective longitudinal studies suggest that engaging in regular physical activity reduces the likelihood of developing a psychiatric illness (Schuch et al., 2018; Suetani et al., 2017), and improves quality of life and psychiatric symptoms in adults with a diagnosable psychiatric illness (Cooney et al., 2013; Rosenbaum et al., 2014). There is also encouraging preliminary evidence that physical activity interventions are effective in the treatment of MDD, with some studies suggesting that physical activity yields symptom improvement comparable to that of treatment as usual (e.g., pharmacotherapy or psychotherapy; Bailey et al., 2018; Davis et al., 2015; Nyström et al., 2015). Therapeutic effects of physical activity have also been found in non-clinical populations (Conn, 2010; Rebar et al., 2015). For instance, a meta-meta-analysis aggregating data from randomized controlled trials (RCTs) of physical activity in community adults observed significant reductions in symptoms of depression and anxiety, with the benefits being most robust for depressive symptoms (Rebar et al., 2015). Physical activity also acutely induces transient elevations in mood (Bernstein and McNally, 2017; Tartar et al., 2018; Zschucke et al., 2015), an effect that emerges even with brief bouts of exercise (e.g., 10 min; Crush et al., 2018). Higher levels of self-reported habitual physical activity have also been associated with lower perceived stress and less aberrant eating in a cross-sectional community sample of pregnant women with overweight and obesity (Donofry et al., 2020a, 2020b), a population at elevated risk for adverse obstetric and maternal health outcomes. Further, among older adults reporting high levels of perceived stress, physical activity engagement has been prospectively associated with reduced ratings of perceived stress and to protect against the development of stress-related physical health symptoms such as chest pain, joint pain, or shortness of breath (Rueggeberg et al., 2012). Together, these studies demonstrate the potential for physical activity interventions to facilitate improvements in psychological health and well-being in both clinical and community samples.

5.2. Cognitive outcomes

Physical activity also has well-documented benefits for cognitive health across the lifespan, demonstrating efficacy for maintenance of cognitive functions as well as for partial remediation of declines in cognitive functioning with illness or aging (Erickson et al., 2019; Groot et al., 2016; Najjar et al., 2019; Oberlin et al., 2017); though see (Brasure et al., 2017; Sabia et al., 2017; Sink et al., 2015) for contradictory patterns. For instance, findings from a meta-analysis of nine prospective studies examining the association between physical activity and risk for AD demonstrated that older adults who engaged in regular physical activity were significantly less likely to develop AD compared to their less active counterparts (Beckett et al., 2015). Physical activity also exerts beneficial effects on cognition among individuals who have

already begun to experience cognitive decline. A meta-analysis of 18 RCTs of physical activity conducted in adults with dementia found that physical activity improved cognitive functioning regardless of the type (aerobic vs. non-aerobic) and frequency of exercise, and the effects did not differ based on dementia type (AD vs. other dementias; (Groot et al., 2016)). In addition to these broad effects on cognition and cognitive decline, there are several sub-domains of cognition that appear to be particularly sensitive to the effects of physical activity. Among older adults, physical activity is most strongly associated with improvements in attention, memory and executive control processes such as working memory, planning, and multi-tasking (Erickson et al., 2019). Findings from several large epidemiological cohort studies of older adults indicate that more frequent engagement in physical activity, particularly of moderate-to-vigorous intensity (MVPA; e.g., brisk walking, jogging), is associated with better performance on tests of delayed memory and verbal fluency (Daly et al., 2015; Zhu et al., 2015). Similar effects have been observed in RCTs of physical activity (Erickson et al., 2011; Groot et al., 2016; Nagamatsu, 2012; Nagamatsu et al., 2013), further strengthening support for implementing physical activity interventions to improve trajectories of cognitive aging. Of note, light intensity activities such as slow paced walking and other such leisure time activities (e.g., light cleaning, gardening) may not promote significant improvements in cognitive performance (Zhu et al., 2015), suggesting that quantity as well as intensity of physical activity behaviors may moderate their impact on cognitive functioning (Lambourne and Tomporowski, 2010; McMorris and Hale, 2012). However, research examining how the dose of physical activity (as measured via quantity and/or intensity) influences cognitive outcomes is relatively limited (Erickson et al., 2019). In addition, comparatively fewer studies have examined the impact of physical activity on cognitive functioning in young and midlife adults, highlighting an important area for future research. Despite these limitations, there is moderate to strong evidence that physical activity enhances cognitive performance and prevents or slows age-related cognitive decline among older adults (Erickson et al., 2019).

5.3. Brain structure and function

In addition to the benefits of physical activity for mood and cognitive functioning, emerging research suggests that physical activity modifies the structure and function of the brain. In particular, physical activity appears to have the most robust effects on prefrontal, parietal and temporal regions that support the same cognitive and psychological processes known to be impaired by ELA. Just as the hippocampus is particularly sensitive to the effects of ELA, the hippocampus appears to be one of the most consistently reported regions affected by physical activity. Numerous studies have demonstrated that aerobic exercise promotes increases in hippocampal volume in children (Migueles et al., 2020) and adults (Erickson et al., 2014, 2011; Firth et al., 2018) and may slow the trajectory of hippocampal volume loss that typically occurs with aging (Erickson et al., 2011; Firth et al., 2018). Furthermore, these effects may emerge relatively rapidly, with one study documenting increases in hippocampal volume and axonal myelination within six weeks of adopting a routine exercise program among midlife adults (Thomas et al., 2016). Importantly, exercise-induced increases in hippocampal volumes have been linked to improvements in cognition. Older adults randomized to 12 months of brisk walking exhibited a two percent increase in hippocampal volume, with increased volume predicting intervention-related improvement in spatial working memory (Erickson et al., 2011). This suggests that the effects of physical activity on brain volume translate to meaningful benefits to other indices of brain health that have a direct impact on quality of life. Physical activity is also associated with the volume of prefrontal cortical regions that support executive processes and typically shrink with advancing age. For instance, MVPA has been positively correlated with dlPFC volume in older adults (Northey et al., 2020). Further, a longitudinal investigation of the effects of regular physical activity found that individuals who

were more active in midlife exhibited higher total prefrontal grey matter volume in late life relative to their sedentary counterparts (Rovio et al., 2010). In addition to these effects on regional grey matter volume, physical activity has also been prospectively associated with reduced deposition of beta-amyloid (A β), a hallmark of AD pathophysiology that develops prior to the onset of cognitive symptoms, among cognitively healthy older adults (Rabin et al., 2019; Stillman et al., 2017). Moreover, physical activity moderated the association between A β accumulation, cognition and brain volume, with more active individuals exhibiting less pronounced declines in both cognitive function and grey matter volume (Rabin et al., 2019). These studies provide evidence that physical activity is protective against age-related volume loss and development of neuropathology characteristic of AD.

Physical activity has also been associated with variation in the function of regions embedded in networks known to be disrupted by ELA (Stillman et al., 2019). Higher levels of physical activity have been positively correlated with DMN connectivity in observational (Boraxbekk et al., 2016; Santaella et al., 2019) and intervention studies (Voss et al., 2010), a pattern which opposes that documented among ELA-exposed adults (e.g., (Sripada et al., 2014)). Further, findings from a recent meta-analysis of 20 exercise RCTs conducted in adults provide converging evidence that exercise is associated with increased activation of regions in the DMN, such as the precuneus and medial temporal lobe (including hippocampus), as well as regions in the executive control network (ECN), including the inferior and superior parietal cortex during cognitively demanding tasks (Yu et al., 2020). Moreover, physical activity was related to increased functional connectivity between sub-regions of the ECN (Yu et al., 2020). Of note, this meta-analysis was limited to studies documenting an association between intervention-related changes in brain activation patterns and changes in cognitive functioning. Thus, these findings provide further evidence that physical activity-induced changes in brain structure and function translate into meaningful improvements in cognitive functioning. Interestingly, a single session of MVPA has been shown to preserve inhibitory control (as measured via performance on the Stroop task) following experimental disruption of dlPFC activity via continuous theta burst stimulation (Lowe et al., 2017). This neuroprotective effect of exercise on inhibitory control was not observed after engagement in light intensity activities (Lowe et al., 2017), consistent with other data suggesting that higher intensity physical activity may be necessary for cognitive benefits to emerge (Lambourne and Tomporowski, 2010; McMorris and Hale, 2012). Together, these studies demonstrate that physical activity promotes neuroplasticity into late adulthood and potentially in the presence of neurodegenerative disease. In light of this evidence, physical activity represents a promising behavioral approach for remediating the deleterious effects of ELA on brain health in adulthood.

6. Overlapping mechanistic pathways through which physical activity might buffer the negative consequences of early life adversity on adult brain health

Given that ELA and physical activity both prompt widespread changes in numerous and overlapping biological systems of the body, physical activity may exert opposing effects on many of the mechanistic pathways through which ELA influences adult brain health. Here, we provide an overview of the literature examining four mechanistic pathways with some of the strongest evidence linked to both ELA and physical activity. These include: (1) neurotrophic factors, (2) HPA axis regulation, (3) inflammatory processes, and (4) epigenetic modifications. Fig. 2 illustrates these pathways and provides a framework for understanding how physical activity may mitigate the negative consequences of ELA on brain health in adulthood. This model is intended to serve as a guide for future research efforts examining the efficacy of physical activity among adults exposed to ELA.




Fig. 2. Theoretical model representing the mechanistic pathways and brain health outcomes associated with early life adversity that are also modified by physical activity. *Note.* BDNF = brain-derived neurotrophic factor; HPA = hypothalamic-pituitary-adrenal.

6.1. Neurotrophic factors

Neurotrophic factors are endogenous peptides that support the survival and growth of developing and mature neurons. They are critically involved in the regulation of experience-dependent brain plasticity. The neurotrophin class of neurotrophic factors have been the most thoroughly investigated for their role in neuronal and synaptic remodeling, particularly brain-derived neurotrophic factor (BDNF), which is abundantly expressed throughout the brain (Thoenen, 1995). Animal models of ELA have shown that exposure to adversities modulates BDNF expression in regions that support learning, memory, cognitive control and emotion regulation, including PFC, hippocampus, and amygdala, and ELA-related disruption of BDNF signaling pathways has been linked to abnormal brain development (Bondar and Merkulova, 2016; Miskolczi et al., 2019). Although human studies have not yet supported a causal link between ELA, BDNF signaling, and brain health, evidence suggests that individuals who are genetically predisposed to reductions in the expression of BDNF are also more vulnerable to the harmful effects of ELA on brain health (Carballedo et al., 2013; Casey et al., 2009; Elzinga et al., 2011).

In contrast, physical activity has been shown to enhance BDNF expression and signaling in both animal and human studies. For instance, experimental studies conducted in rodents have shown that aerobic exercise increases BDNF levels in the striatum (Marais et al., 2009), hippocampus (Vaynman et al., 2004; Wrann et al., 2013), and PFC (D.-C. Wang et al., 2020). Moreover, the neuroprotective effects of aerobic exercise in animal models of Parkinson's Disease and AD are dependent in part on increased production of BDNF (Real et al., 2013; Xiong et al., 2015), with blockade of BDNF receptors attenuating exercise-induced improvements in brain health (Real et al., 2013; Vaynman et al., 2004). Observational studies in humans have demonstrated that both acute and habitual aerobic exercise are associated with increases in peripheral BDNF levels (Dinoff et al., 2016; Szuhany et al., 2015). Further, higher concentrations of peripheral BDNF following participation in an exercise intervention mediated intervention-related improvements in executive functioning (Leckie et al., 2014) and memory (Whiteman et al., 2014). Exercise-induced increases in serum BDNF levels have also been correlated with increased hippocampal volume (Erickson et al., 2011) and functional connectivity (Voss et al., 2013) following an exercise intervention, providing additional support for the hypothesis that exercise promotes brain health through modulation of BDNF signaling pathways. As such, it is possible that engaging in physical activity may mitigate the effects of ELA on brain health by promoting BDNF expression.

6.2. HPA axis regulation

The HPA axis is the principle neuroendocrine system responsible for

orchestrating physiological responses to physical or psychological challenges, with these responses being mediated by the systemic release of the glucocorticoid hormone cortisol from the adrenal cortex (Spencer and Deak, 2017). Prolonged exposure to events that stimulate HPA axis reactivity, such as is the case with ELA, disrupts homeostatic regulation of this system, thus reducing the capacity of the body to adequately respond to new or worsening challenges (Jankord and Herman, 2008). Indeed, ELA disrupts the development of the HPA axis by programming the system to respond sub-optimally to environmental demands (McLaughlin et al., 2015; van Bodegom et al., 2017). The impact of ELA on HPA axis regulation is complex and may vary across developmental periods or based on the type, timing, and severity of ELA exposure, as well as according to whether studies are examining basal or stressor-evoked HPA axis activation (Bunea et al., 2017; van Bodegom et al., 2017). Accordingly, some studies have observed hyper-reactivity of the HPA axis among adults exposed to ELA (Butler et al., 2017; Friedman et al., 2015), while others have observed blunting of HPA axis activation (Bunea et al., 2017; Karlamangla et al., 2019; Kuras et al., 2017; Lovallo et al., 2013). Nevertheless, there is general scientific consensus that childhood adversity is associated with altered HPA functioning in adulthood. Of note, the pro-inflammatory state provoked by ELA has been implicated in abnormal development of the HPA axis (Nusslock and Miller, 2016), suggesting that dysregulation of one biological system following exposure to ELA may potentiate dysregulation in other systems involved in maintaining brain health. Further, disruption of the HPA axis has also been associated with increased risk for psychiatric and medical conditions that are prevalent among adults with a history of ELA, including MDD (Belvederi Murri et al., 2014), Type II diabetes (Joseph and Golden, 2017), cardiovascular disease (Jokinen and Nordström, 2009), and AD (Popp et al., 2015). Moreover, several brain regions involved in cognitive and emotional processing, most notably the hippocampus, densely express glucocorticoid receptors and are therefore sensitive to maladaptation in the HPA system (Gianaros et al., 2007; Meijer et al., 2019; Seo et al., 2019). Thus, dysregulation of the HPA axis following exposure to ELA is likely a mechanism underlying heightened vulnerability to stress-related psychopathology and chronic disease.

Physical activity behaviors are energetically demanding and therefore require robust neuroendocrine adaptations in order to mobilize resources and maintain physiological homeostasis. As such, it is perhaps unsurprising that physical activity evokes acute changes in HPA axis activation that mirror those observed in the context of psychosocial stress. However, according to the cross-stressor adaptation hypothesis (Sothmann et al., 1996), habitual engagement in physical activity may facilitate homeostatic regulation of the HPA system and mitigate disruption of HPA axis functioning prompted by chronic exposure to psychosocial stressors, including ELA (Zschucke et al., 2015). Individuals who engage in regular physical activity exhibit attenuated

cortisol responses to laboratory-based mental stress tasks (Gerber et al., 2017; Rimmele et al., 2009; Strahler et al., 2016; Wood et al., 2017; Wunsch et al., 2019), including those reporting elevated negative affect (Pauly et al., 2019; Puterman et al., 2011). Although there have been relatively few RCTs of physical activity to examine change in HPA axis regulation as an outcome of intervention, there is some preliminary data to suggest that habitual physical activity alters HPA axis functioning (Drogos et al., 2019; Klaperski et al., 2014; Saxton et al., 2014), converging with evidence from the aforementioned cross-sectional studies. Interestingly, regular engagement in mind-body interventions that combine elements of mindfulness and physical movement practices such as tai-chi and yoga have also been shown to reduce basal cortisol output (Campo et al., 2015; Gothe et al., 2016; Sarubin et al., 2014). For instance, older adults randomized to an eight-week hatha yoga program exhibited reduced cortisol levels following the intervention, an effect that was not observed in a stretching control group (Gothé et al., 2016). Further, reductions in cortisol were associated with intervention-related improvements in executive functioning (Gothé et al., 2016). Together, these data suggest that physical activity may promote brain health by improving HPA axis regulation, which may help to remediate the enduring physiological impacts of childhood adversity.

6.3. Inflammatory processes

Dysregulation of the immune system is one pathway through which ELA may become biologically embedded and influence brain health across the lifespan (Muscatell et al., 2020; Nusslock and Miller, 2016). Exposure to ELA has been associated with chronic elevation in circulating pro-inflammatory molecules such as C-reactive protein (CRP) and cytokines such as interleukin-6 (IL-6) and tumor necrosis factor – alpha (TNF- α), with this pro-inflammatory state persisting into adulthood (Baumeister et al., 2016). For instance, a small ($N = 69$) study in which participants were exposed to an acute stressor found evidence that adults with a self-reported history of childhood maltreatment exhibited greater acute IL-6 release following stressor exposure, as well as higher sustained IL-6 concentrations, compared to individuals without a history of maltreatment (Carpenter et al., 2010). Interestingly, all participants were free of psychiatric diagnoses such as major depression or post-traumatic stress disorder, suggesting that group differences in IL-6 concentrations were not attributable to a higher prevalence of these conditions in adults with a history of maltreatment. Data drawn from the MIDUS and 1958 British cohorts ($N = 8916$) similarly found that exposure to ELA, particularly physical abuse, was associated with elevations of basal CRP and fibrinogen, a glycoprotein that facilitates blood clotting (Pinto Pereira et al., 2019). These associations were attenuated when accounting for adult adiposity, indicating that obesity may partially mediate the relationship between ELA and adult inflammation (Pinto Pereira et al., 2019). Of note, Black race has been shown to prospectively predict higher levels of circulating CRP and hypertension in adulthood independently of other factors such as family and neighborhood poverty measured in childhood (Nikulina and Widom, 2014). These startling findings suggest that race and racism are independently related to poor health in adulthood above and beyond childhood socioeconomic indicators known to robustly predict poor health outcomes. Importantly, prolonged dysregulation of the immune system such as that observed among individuals with a history of ELA may in turn increase vulnerability for numerous psychiatric and medical conditions in adulthood, including MDD (Stewart et al., 2009; Zalli et al., 2016), cardiovascular disease (Packard et al., 2011; Ruparelia et al., 2017; Stewart et al., 2009), and mild cognitive impairment and dementia (Darweesh et al., 2018; Ozawa et al., 2017). Further, elevations in circulating IL-6 have been shown to mediate the relationship between childhood adversity and functional connectivity between regions in the corticolimbic circuit that support affect regulation (Kraynak et al., 2019). Thus, ELA may contribute to poor brain health in adulthood by provoking chronic pro-inflammatory processes, potentially initiating or

exacerbating a cascade of negative physical and psychological health outcomes.

Habitual physical activity, conversely, has been associated with improved inflammatory profiles. Large observational studies have demonstrated that both self-reported (Mayat et al., 2020; Papenberg et al., 2016; Vella et al., 2017) and objectively measured physical activity (Draganidis et al., 2018; Nilsson et al., 2018) are negatively correlated with circulating levels of pro-inflammatory markers such as IL-6 and CRP, even after adjusting for factors such as central adiposity known to promote inflammation (Hamer et al., 2012; Vella et al., 2017). Interestingly, data from a large epidemiological study found evidence that the frequency (i.e., number of days per week) of objectively measured MVPA was a stronger predictor of circulating CRP than was total volume (i.e., minutes per week) among adults (Loprinzi, 2015), suggesting that consistent engagement in MVPA may be necessary to elicit anti-inflammatory effects. However, because few studies have compared the impact of various parameters of physical activity on such outcomes, additional research is needed to strengthen this hypothesis. Prospective cohort studies have similarly found that higher habitual physical activity at baseline, particularly MVPA, predicts lower concentrations of IL-6 and CRP up to 10 years later (Braskie et al., 2014; Fuentes et al., 2020; Hamer et al., 2012; Martinez-Gomez et al., 2019; Menezes et al., 2019). Data from randomized controlled trials (RCT) of physical activity lasting between four and 78 weeks also indicate that structured physical activity programs reduce inflammation among healthy adults as well as in adults with chronic medical conditions (Freitas et al., 2017; Lin et al., 2015; Tartibian et al., 2015; Tomeleri et al., 2016), providing more definitive evidence that physical activity improves immune system functioning. Interestingly, the pro-inflammatory effects of cancer treatments such as chemotherapy and radiation therapy may be mitigated by physical activity. Several intervention trials conducted among women with breast cancer found that women randomized to a physical activity program did not exhibit elevations in circulating inflammatory markers that typically occur following cancer treatment, while women in the control groups demonstrated the expected post-treatment spike in cytokine concentrations (Hagstrom et al., 2016; Mustian et al., 2015; Schmidt et al., 2016). Together, these studies suggest that physical activity reduces inflammation and may therefore be an effective approach to restoring immune system functioning following exposure to ELA.

6.4. Epigenetic modifications

Gene expression is regulated not only by transcription and translation of DNA, but also through experience-dependent modification of the structure of chromatin, which determines the degree to which a transcription site is exposed and available to initiate gene transcription (Jaenisch and Bird, 2003). DNA Methylation, the process through which methyl groups are added to the cytosine nucleotide, is the most common epigenetic mechanism through which environmental exposures alter the expression potential of genes (Jaenisch and Bird, 2003). An emerging body of research suggests that ELA induces widespread epigenetic modifications that may underlie many of the observed physiological, behavioral, and neural abnormalities in adulthood that have been linked to ELA (Vaiserman and Koliada, 2017). A study of post-mortem brain tissue among individuals who completed suicide identified 362 promoter regions at which the degree of methylation differed between those with and without a history of ELA, with many of the affected genes being involved in neural development and plasticity (Labonté et al., 2012). Famine exposure in infancy has been associated with hypermethylation of genes that regulate energy and lipid metabolism in adulthood, and the degree of methylation in these genes mediated the relationship between famine exposure and adult body composition and circulating triglyceride levels (Tobi et al., 2018), suggesting that the impact of ELA on gene expression patterns is enduring and may influence key indicators of health in adulthood. Longitudinal analyses in the

New England Family Study showed evidence that several methylation sites in adipose tissue mediated the relation between childhood socioeconomic adversity and adulthood adiposity, including many sites that are biologically relevant for the development of obesity, such as fatty acid synthase, transmembrane protein 88, and neuritin 1 (Loucks et al., 2016). Of particular relevance to brain health, methylation of the BDNF gene has been shown to be elevated in adults reporting ELA (J. A. Smith et al., 2017; Unternaehrer et al., 2015), which may reduce neural plasticity in adulthood and contribute to premature declines in brain health.

Other studies have demonstrated that genes involved in HPA axis and immune system regulation are particularly sensitive to ELA. For instance, childhood adversity has been associated with hypermethylation of the gene encoding the glucocorticoid receptor, NR3C1 (Farrell et al., 2018; Tyrka et al., 2012) and the FKBP5 gene, which encodes a protein that modulates the sensitivity of the glucocorticoid receptor (Harms et al., 2017; Klengel et al., 2013). Further, FKBP5 hypermethylation negatively correlated with dorsolateral PFC activation during the performance of a Go/No-go task among adults with a history of ELA (Harms et al., 2017), providing preliminary evidence that epigenetic modifications affecting HPA axis regulation may disrupt neurocognitive functioning. Methylation of immune-regulating genes has also been associated with ELA, with one study demonstrating that neighborhood disadvantage in childhood related to hypermethylation of seven genes involved in inflammatory processes (J. A. Smith et al., 2017). In addition, methylation of a IL-6 promoter region has been shown to be reduced among individuals exposed to neighborhood violence and other more severe forms of childhood trauma, and predicted elevated levels of circulating IL-6 evoked by a psychosocial stressor (Janusek et al., 2017). Thus, exposure to childhood adversity may become biologically embedded through epigenetic modifications to genes that regulate processes integral to neurodevelopment and the maintenance of brain health across the lifespan. Given that research examining the impact of ELA on DNA methylation in humans has only emerged relatively recently, there is a need for additional well-designed studies linking ELA-related epigenetic modifications to brain health outcomes in adulthood.

Physical activity is similarly associated with widespread epigenetic modifications that may underlie its observed benefits for health and well-being. Overall, DNA methylation across the genome is positively correlated with the extent of routine physical activity engagement, though the consequences of physical activity-related methylation patterns are varied and dependent upon the molecular pathways affected by changes in gene expression (Boyne et al., 2018; F. F. Zhang et al., 2011). For instance, habitually engaging in physical activity of any intensity has been associated with higher methylation of the apoptosis associated speck-like protein containing a caspase recruitment domain (ASC) gene, which encodes a protein involved in cellular apoptosis and stimulation of IL-1 β cytokine release (Nakajima et al., 2010; Nishida et al., 2019). Further, observational evidence suggests that ASC methylation is positively correlated with aerobic fitness capacity and mediates the effect of cardiorespiratory fitness on reductions in IL-1 β levels (Butts et al., 2017), indicating that increased ASC methylation due to routine physical activity may promote a more favorable immune profile. Engaging in three months of physical activity has also been related to increased methylation of genes that encode proteins that regulate endothelial function and vasoconstriction, including Endothelin-1 and Nitrous Oxide Synthase-2, with increased methylation of these genes predicting improvements in blood pressure (Ferrari et al., 2019). In contrast, data from an RCT of aerobic exercise among women with breast cancer demonstrated that physical activity reduced methylation of a tumor suppressor gene, L3MBTL1, thus increasing expression of a protein that mitigated tumor progression (Zeng et al., 2012). Importantly, physical activity-induced reductions in L3MBTL1 methylation predicted lower rates of breast cancer recurrence and mortality (Zeng et al., 2012). These findings indicate that physical activity influences health through

epigenetic modifications of many of the same molecular pathways impacted by ELA.

7. Summary and future directions

Early life adversity exerts broad, profound and enduring effects on brain health in adulthood. Evidence suggests that physical activity interventions also affect multiple indices of brain health and modify the molecular pathways through which ELA impacts brain health, including neurotrophic factors, HPA axis regulation, inflammatory processes, and epigenetic mechanisms. However, although physical activity interventions show promise as a low-cost behavioral approach to enhancing brain health, it remains unclear whether the benefits of physical activity extend to individuals who have been exposed to ELA. There are a number of important areas for future investigation that will serve to establish the efficacy of physical activity for remediating the deleterious effects of ELA on brain health. As an initial step toward this aim, it will be necessary to conduct observational studies exploring whether ELA moderates the relationship between physical activity and brain health outcomes in adulthood, both cross-sectionally and longitudinally. It will also be critical to conduct rigorous RCTs of physical activity to examine whether ELA impacts the efficacy of physical activity interventions, and if so, the mechanisms underlying ELA-related differences in outcomes. For instance, adverse childhood experiences increase vulnerability to adversities in adulthood, including lower socioeconomic status and higher risk of work disability (Halonen et al., 2017; Montez and Hayward, 2014). The socioeconomic and socioemotional consequences of ELA are significant barriers to participation in research, particularly time intensive clinical trials of physical activity that may require numerous in-person visits each week over the course of several months. Therefore, to better serve this population, the format in which physical activity is delivered may need to be adapted to accommodate the challenges associated with ELA. Modifications to the type (e.g., aerobic versus resistance), intensity, format (e.g., in-person versus virtual), and duration of physical activity interventions should be explored, followed by comparative investigation of the effectiveness of tailored versus standardized interventions. This should include exploration of the impact of physical activities not typically well-represented in RCTs, including dance, yoga, martial arts, and structured sports programs. Relatedly, it is likely that the type, timing, and severity of ELA will impact intervention outcomes. Thus, careful and thorough evaluation of ELA histories will be crucial for establishing whether individual differences in ELA exposure profiles moderate intervention effects. Given the aforementioned issues that remain regarding how to best operationalize ELA and its neurodevelopmental consequences, it will be important for research in this area to be guided by a sound theoretical framework with explicit disclosure of the assumptions and limitations of the chosen framework.

Additional research is also needed to determine whether the proposed mechanistic pathways through which physical activity may promote brain health are indeed related to intervention outcomes among adults exposed to ELA, and to compare the extent to which each pathway mediates intervention effects. It is also possible that certain brain health outcomes are more responsive to intervention than others, highlighting the need for studies that measure multiple indices of brain health to evaluate whether there is specificity of the effects of physical activity. It will also be important to compare the effects of physical activity among adults exposed to ELA to outcomes observed in existing psychotherapeutic approaches to treat trauma, such as trauma-focused Cognitive Behavioral Therapy, Prolonged Exposure Therapy, or Cognitive Processing Therapy. Finally, given evidence that early intervention typically produces more robust and durable improvements (Karoly et al., 2005), it will be important to examine whether the age at the time of the intervention delivery moderates the efficacy of physical activity for remediating the negative health consequences of ELA. It is clear that physical activity is effective for enhancing brain health in late

adulthood, even among individuals who have begun to experience neurocognitive decline (Chirles et al., 2017; Hillman et al., 2008; Oberlin et al., 2017). However, it may be the case that older age in combination with a history of ELA limits the impact of physical activity on indicators of brain health. Alternatively, should physical activity prove to be effective in this population, it would suggest that there is not a “point of no return” beyond which the negative consequences of ELA become irreversible.

Finally, there is a critical need for research exploring the impact of racism and other forms of discrimination on brain health in adulthood, and to determine whether and how physical activity may mitigate these effects. It is clear that there are marked racial disparities in the prevalence of age-related cognitive impairment and neurocognitive illnesses such as AD, with Black and African American individuals being at two to three times higher risk for such outcomes compared to White individuals (Lee et al., 2012; Mehta and Yeo, 2017; Weuve et al., 2018). These disparities are driven in part by the fact that psychiatric and cardiometabolic risk factors for cognitive decline and dementia are more prevalent in minority communities (Eack and Newhill, 2012; McGuire and Miranda, 2008; Topel et al., 2018; Wassink et al., 2017). However, given that minority communities are also disproportionately burdened by ELA of all forms (Merrick et al., 2018), exposure to ELA is likely to account for some proportion of racial health disparities. Further, minority communities experience the additional burden of racism, a form of adversity known to uniquely impact adult health outcomes (Nikulina and Widom, 2014). This has prompted calls to expand theoretical models of ELA to include racism as an independent form of ELA to promote research aimed at better understanding the influence of racism-related experiences on health and well-being across the lifespan (Bernard et al., 2020). Yet, there is a relative paucity of research to specifically explore the contribution of racism experienced during childhood and adolescence to the emergence of racial disparities in brain health in mid- and late life. The impact of racism on physical activity engagement and on the efficacy of physical activity interventions for improving aspects of brain health is also understudied. These represent critical scientific gaps that must be addressed in order to narrow racial disparities in brain health.

To conclude, physical activity represents a promising, low-cost and readily disseminated approach to mitigating or remediating the negative consequences of ELA on brain health in adulthood. However, given the lack of studies exploring the efficacy of physical activity for improving brain health outcomes specifically among adults affected by ELA, additional research is needed to evaluate the hypotheses put forth in this review. Pursuit of the research aims outlined above will establish whether physical activity modifies the processes that underlie the long-term consequences of ELA on brain health, which has the potential to benefit the millions of adults globally who have been exposed to ELA.

Sources of funding

This manuscript was supported by grants from the National Institute of Aging (R24 AG065174 [PI: E. Loucks, M. Sheridan, and K. Godfrey]; R01 AG060741 [PI: K. Erickson]), the National Heart Lung and Blood Institute (T32 HL007560 [PI: K. Matthews; awardee: S. Donofry]), and the National Center for Complementary and Integrative Health (K23 AT011173 [PI: S. Sun]). The funding sources had no involvement in the conceptualization, writing of the manuscript, or in the decision to submit the manuscript for publication.

Declaration of Competing Interest

The authors have no conflicts of interest to disclose.

Acknowledgements

We would like to thank Alysha Gilmore, a research assistant in the

Brain Aging and Cognitive Health laboratory for her assistance with organizing the literature review for this manuscript.

References

- Acheson, A., Wijtenburg, S.A., Rowland, L.M., Winkler, A.M., Gaston, F., Mathias, C.W., et al., 2014. Assessment of whole brain white matter integrity in youths and young adults with a family history of substance-use disorders. *Hum. Brain Mapp.* 35 (11), 5401–5413. <https://doi.org/10.1002/hbm.22559>.
- Almanza-Sepulveda, M.L., Chico, E., Gonzalez, A., Hall, G.B., Steiner, M., Fleming, A.S., 2018. Executive function in teen and adult women: association with maternal status and early adversity. *Dev. Psychobiol.* 60 (7), 849–861. <https://doi.org/10.1002/dev.21766>.
- Bailey, A.P., Hetrick, S.E., Rosenbaum, S., Purcell, R., Parker, A.G., 2018. Treating depression with physical activity in adolescents and young adults: a systematic review and meta-analysis of randomised controlled trials. *Psychol. Med.* 48 (7), 1068–1083. <https://doi.org/10.1017/S0033291717002653>.
- Banihashemi, L., Sheu, L.K., Midei, A.J., Gianaros, P.J., 2015. Childhood physical abuse predicts stressor-evoked activity within central visceral control regions. *Soc. Cogn. Affect. Neurosci.* 10 (4), 474–485. <https://doi.org/10.1093/scan/nsu073>.
- Baumeister, D., Akhtar, R., Ciufolini, S., Pariante, C.M., Mondelli, V., 2016. Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor- α . *Mol. Psychiatry* 21 (5), 642–649. <https://doi.org/10.1038/mp.2015.67>.
- Beck, A., Franz, C.E., Xian, H., Vuoksimaa, E., Tu, X., Reynolds, C.A., et al., 2018. Mediators of the effect of childhood socioeconomic status on late midlife cognitive abilities: a four decade longitudinal study. *Innov. Aging* 2 (1). <https://doi.org/10.1093/geroni/igy003>.
- Beckett, M.W., Arden, C.L., Rotondi, M.A., 2015. A meta-analysis of prospective studies on the role of physical activity and the prevention of Alzheimer's disease in older adults. *BMC Geriatr.* 15 (1), 9. <https://doi.org/10.1186/s12877-015-0007-2>.
- Belsky, D.W., Caspi, A., Houts, R., Cohen, H.J., Corcoran, D.L., Danese, A., et al., 2015. Quantification of biological aging in young adults. *Proc. Natl. Acad. Sci.* 112 (30), E4104–E4110. <https://doi.org/10.1073/pnas.1506264112>.
- Belvederi Murri, M., Pariante, C., Mondelli, V., Masotti, M., Atti, A.R., Mellacqua, Z., et al., 2014. HPA axis and aging in depression: systematic review and meta-analysis. *Psychoneuroendocrinology* 41, 46–62. <https://doi.org/10.1016/j.psyneuen.2013.12.004>.
- Bernard, D.L., Calhoun, C.D., Banks, D.E., Halliday, C.A., Hughes-Halbert, C., Danielson, C.K., 2020. Making the “C-ACE” for a culturally-informed adverse childhood experiences framework to understand the pervasive mental health impact of racism on black youth. *J. Child Adolesc. Trauma.* <https://doi.org/10.1007/s40653-020-00319-9>.
- Bernstein, E.E., McNally, R.J., 2017. Acute aerobic exercise helps overcome emotion regulation deficits. *Cogn. Emot.* 31 (4), 834–843. <https://doi.org/10.1080/02699931.2016.1168284>.
- Bernstein, E.E., Heeren, A., McNally, R.J., 2017. Unpacking rumination and executive control: a network perspective. *Clin. Psychol. Sci.* 5 (5), 816–826. <https://doi.org/10.1177/2167702617702717>.
- Bick, J., Zhu, T., Stamoulis, C., Fox, N.A., Zeanah, C., Nelson, C.A., 2015. Effect of early institutionalization and foster care on long-term white matter development: a randomized clinical trial. *JAMA Pediatr.* 169 (3), 211–219. <https://doi.org/10.1001/jamapediatrics.2014.3212>.
- Birn, R.M., Roeber, B.J., Pollak, S.D., 2017. Early childhood stress exposure, reward pathways, and adult decision making. *Proc. Natl. Acad. Sci.* 114 (51), 13549–13554. <https://doi.org/10.1073/pnas.1708791114>.
- Boecker, R., Holz, N.E., Buchmann, A.F., Blomeyer, D., Plichta, M.M., Wolf, I., et al., 2014. Impact of early life adversity on reward processing in young adults: EEG-fMRI results from a prospective study over 25 years. *PLoS One* 9 (8), e104185. <https://doi.org/10.1371/journal.pone.0104185>.
- Boecker-Schlier, R., Holz, N.E., Buchmann, A.F., Blomeyer, D., Plichta, M.M., Jennen-Steinmetz, C., et al., 2016. Interaction between COMT Val158Met polymorphism and childhood adversity affects reward processing in adulthood. *NeuroImage* 132, 556–570. <https://doi.org/10.1016/j.neuroimage.2016.02.006>.
- Bondar, N.P., Merkulova, T.L., 2016. Brain-derived neurotrophic factor and early-life stress: multifaceted interplay. *J. Biosci.* 41 (4), 751–758. <https://doi.org/10.1007/s12038-016-9648-3>.
- Boots, E.A., Schultz, S.A., Oh, J.M., Larson, J., Edwards, D., Cook, D., et al., 2015. Cardiorespiratory fitness is associated with brain structure, cognition, and mood in a middle-aged cohort at risk for Alzheimer's disease. *Brain Imaging Behav.* 9 (3), 639–649. <https://doi.org/10.1007/s11682-014-9325-9>.
- Boraxbekk, C.-J., Salami, A., Wåhlin, A., Nyberg, L., 2016. Physical activity over a decade modifies age-related decline in perfusion, gray matter volume, and functional connectivity of the posterior default-mode network—a multimodal approach. *NeuroImage* 131, 133–141. <https://doi.org/10.1016/j.neuroimage.2015.12.010>.
- Botros, M., Hodgins, G., Nemeroff, C.B., 2019. Chapter 11 - the Long-lasting neurobiological scars of early-life stress: implications for the neurobiology of depression. In: Quevedo, J., Carvalho, A.F., Zarate, C.A. (Eds.), *Neurobiology of Depression*. Academic Press, pp. 111–121. <https://doi.org/10.1016/B978-0-12-813333-0.00011-1>.
- Boyne, D.J., O'Sullivan, D.E., Olij, B.F., King, W.D., Friedenreich, C.M., Brenner, D.R., 2018. Physical activity, global DNA methylation, and breast Cancer risk: a systematic literature review and meta-analysis. *Cancer Epidemiology and Prevention Biomarkers* 27 (11), 1320–1331. <https://doi.org/10.1158/1055-9965.EPI-18-0175>.

- Braskie, M.N., Boyle, C.P., Rajagopalan, P., Gutman, B.A., Toga, A.W., Raji, C.A., et al., 2014. Physical activity, inflammation, and volume of the aging brain. *Neuroscience* 273, 199–209. <https://doi.org/10.1016/j.neuroscience.2014.05.005>.
- Brasler, M., Desai, P., Davila, H., Nelson, V.A., Calvert, C., Jutkowitz, E., et al., 2017. Physical activity interventions in preventing cognitive decline and alzheimer-type dementia. *Ann. Intern. Med.* 168 (1), 30–38. <https://doi.org/10.7326/M17-1528>.
- Brooks, S.J., Naidoo, V., Roos, A., Fouché, J.-P., Lochner, C., Stein, D.J., 2016. Early-life adversity and orbitofrontal and cerebellar volumes in adults with obsessive-compulsive disorder: voxel-based morphometry study. *The British Journal of Psychiatry: The Journal of Mental Science* 208 (1), 34–41. <https://doi.org/10.1192/bjp.bp.114.162610>.
- Bruni, A., Carbone, E.A., Pugliese, V., Aloï, M., Calabrò, G., Cerminara, G., et al., 2018. Childhood adversities are different in schizophrenic Spectrum disorders, bipolar disorder and major depressive disorder. *BMC Psychiatry* 18 (1), 391. <https://doi.org/10.1186/s12888-018-1972-8>.
- Buckner, R.L., DiNicola, L.M., 2019. The brain's default network: updated anatomy, physiology and evolving insights. *Nat. Rev. Neurosci.* 20 (10), 593–608. <https://doi.org/10.1038/s41583-019-0212-7>.
- Bunea, I.M., Szentágotai-Tátar, A., Miu, A.C., 2017. Early-life adversity and cortisol response to social stress: a meta-analysis. *Transl. Psychiatry* 7 (12), 1–8. <https://doi.org/10.1038/s41398-017-0032-3>.
- Butler, K., Klaus, K., Edwards, L., Pennington, K., 2017. Elevated cortisol awakening response associated with early life stress and impaired executive function in healthy adult males. *Horm. Behav.* 95, 13–21. <https://doi.org/10.1016/j.yhbeh.2017.07.013>.
- Butts, B., Butler, J., Dunbar, S.B., Corwin, E., Gary, R.A., 2017. ASC methylation and Interleukin-1 β are associated with aerobic capacity in heart failure. *Med. Sci. Sports Exerc.* 49 (6), 1072–1078. <https://doi.org/10.1249/MSS.0000000000001200>.
- Byers, A.L., Yaffe, K., 2011. Depression and risk of developing dementia. *Nat. Rev. Neurol.* 7 (6), 323–331. <https://doi.org/10.1038/nrneurol.2011.60>.
- Cameron, L.D., Carroll, P., Hamilton, W.K., 2018. Evaluation of an intervention promoting emotion regulation skills for adults with persisting distress due to adverse childhood experiences. *Child Abuse Negl.* 79, 423–433. <https://doi.org/10.1016/j.chiabu.2018.03.002>.
- Campo, R.A., Light, K.C., O'Connor, K., Nakamura, Y., Lipschitz, D., LaStayo, P.C., et al., 2015. Blood pressure, salivary cortisol, and inflammatory cytokine outcomes in senior female cancer survivors enrolled in a tai chi chih randomized controlled trial. *J. Cancer Surviv.* 9 (1), 115–125. <https://doi.org/10.1007/s11764-014-0395-x>.
- Carballedo, A., Morris, D., Zill, P., Fahey, C., Reinhold, E., Meisenzahl, E., et al., 2013. Brain-derived neurotrophic factor Val66Met polymorphism and early life adversity affect hippocampal volume. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* 162 (2), 183–190. <https://doi.org/10.1002/ajmg.b.32130>.
- Carbone, E.A., Pugliese, V., Bruni, A., Aloï, M., Calabrò, G., Jaén-Moreno, M.J., et al., 2019. Adverse childhood experiences and clinical severity in bipolar disorder and schizophrenia: a transdiagnostic two-step cluster analysis. *J. Affect. Disord.* 259, 104–111. <https://doi.org/10.1016/j.jad.2019.08.049>.
- Carpenter, L.L., Gawuga, C.E., Tyrka, A.R., Lee, J.K., Anderson, G.M., Price, L.H., 2010. Association between plasma IL-6 response to acute stress and early-life adversity in healthy adults. *Neuropsychopharmacology* 35 (13), 2617–2623. <https://doi.org/10.1038/npp.2010.159>.
- Casey, B.J., Glatt, C.E., Tottenham, N., Soliman, F., Bath, K., Amso, D., et al., 2009. Brain-derived neurotrophic factor as a model system for examining gene by environment interactions across development. *Neuroscience* 164 (1), 108–120. <https://doi.org/10.1016/j.neuroscience.2009.03.081>.
- Chirles, T.J., Reiter, K., Weiss, L.R., Alfini, A.J., Nielson, K.A., Smith, J.C., 2017. Exercise training and functional connectivity changes in mild cognitive impairment and healthy elders. *Journal of Alzheimer's Disease: JAD* 57 (3), 845–856. <https://doi.org/10.3233/JAD-161151>.
- Conn, V.S., 2010. Depressive symptom outcomes of physical activity interventions: meta-analysis findings. *Ann. Behav. Med.* 39 (2), 128–138. <https://doi.org/10.1007/s12160-010-9172-x>.
- Cooney, G.M., Dwan, K., Greig, C.A., Lawlor, D.A., Rimer, J., Waugh, F.R., et al., 2013. Exercise for depression. *Cochrane Database Syst. Rev.* 9. <https://doi.org/10.1002/14651858.CD004366.pub6>.
- Crush, E.A., Frith, E., Loprinzi, P.D., 2018. Experimental effects of acute exercise duration and exercise recovery on mood state. *J. Affect. Disord.* 229, 282–287. <https://doi.org/10.1016/j.jad.2017.12.092>.
- Currie, C.L., Sanders, J.L., Swanepoel, L.-M., Davies, C.M., 2020. Maternal adverse childhood experiences are associated with binge drinking during pregnancy in a dose-dependent pattern: findings from the All our Families cohort. *Child Abuse Negl.* 101, 104348. <https://doi.org/10.1016/j.chiabu.2019.104348>.
- Daly, M., McMinn, D., Allan, J.L., 2015. A bidirectional relationship between physical activity and executive function in older adults. *Front. Hum. Neurosci.* 8. <https://doi.org/10.3389/fnhum.2014.01044>.
- Danese, A., McEwen, B.S., 2012. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol. Behav.* 106 (1), 29–39. <https://doi.org/10.1016/j.physbeh.2011.08.019>.
- Dannlowski, U., Stuhrmann, A., Beutelmann, V., Zwanzger, P., Lenzen, T., Grotegerd, D., et al., 2012. Limbic scars: long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biol. Psychiatry* 71 (4), 286–293. <https://doi.org/10.1016/j.biopsych.2011.10.021>.
- Darweesh, S.K.L., Wolters, F.J., Ikram, M.A., de Wolf, F., Bos, D., Hofman, A., 2018. Inflammatory markers and the risk of dementia and Alzheimer's disease: a meta-analysis. *Alzheimers Dementia* 14 (11), 1450–1459. <https://doi.org/10.1016/j.jalz.2018.02.014>.
- Davis, K., Goodman, S.H., Leiferman, J., Taylor, M., Dimidjian, S., 2015. A randomized controlled trial of yoga for pregnant women with symptoms of depression and anxiety. *Complement. Ther. Clin. Pract.* 21 (3), 166–172. <https://doi.org/10.1016/j.ctcp.2015.06.005>.
- DeDonno, S.R., Mickey, B.J., Pruitt, P.J., Stange, J.P., Hsu, D.T., Weldon, A.L., et al., 2019. Influence of childhood adversity, approach motivation traits, and depression on individual differences in brain activation during reward anticipation. *Biol. Psychol.* 146, 107709. <https://doi.org/10.1016/j.biopsycho.2019.05.009>.
- Dinoff, A., Herrmann, N., Swardfager, W., Liu, C.S., Sherman, C., Chan, S., Lancôt, K.L., 2016. The effect of exercise training on resting concentrations of peripheral brain-derived neurotrophic factor (BDNF): a meta-analysis. *PLoS One* 11 (9), e0163037. <https://doi.org/10.1371/journal.pone.0163037>.
- Donofry, S.D., Germeroth, L.J., Kolko Conlon, R.P., Venditti, E.M., Levine, M.D., 2020a. Correlates of physical activity engagement among pregnant women with overweight and obesity. *Women Health Issues* 30 (5), 393–400. <https://doi.org/10.1016/j.whi.2020.06.001>.
- Donofry, S.D., Jakicic, J.M., Rogers, R.J., Watt, J.C., Roeklein, K.A., Erickson, K.I., 2020b. Comparison of food cue-evoked and resting-state functional connectivity in obesity. *Psychosom. Med.* 82 (3), 261–271. <https://doi.org/10.1097/PSY.0000000000000769>.
- Draganidis, D., Jamurtas, A.Z., Stampoulis, T., Laschou, V.C., Deli, C.K., Georgakouli, K., et al., 2018. Disparate habitual physical activity and dietary intake profiles of elderly men with low and elevated systemic inflammation. *Nutrients* 10 (5). <https://doi.org/10.3390/nu10050566>.
- Drogos, L.L., Wynne-Edwards, K., Zhou, R., Hall, S.E., Tyndall, A.V., Longman, R.S., et al., 2019. Aerobic exercise increases cortisol awakening response in older adults. *Psychoneuroendocrinology* 103, 241–248. <https://doi.org/10.1016/j.psyneuen.2019.01.012>.
- Durstine, J.L., Gordon, B., Wang, Z., Luo, X., 2013. Chronic disease and the link to physical activity. *J. Sport Health Sci.* 2 (1), 3–11. <https://doi.org/10.1016/j.jshs.2012.07.009>.
- Duval, E.R., Garfinkel, S.N., Swain, J.E., Evans, G.W., Blackburn, E.K., Angstadt, M., et al., 2017. Childhood poverty is associated with altered hippocampal function and visuospatial memory in adulthood. *Dev. Cogn. Neurosci.* 23, 39–44. <https://doi.org/10.1016/j.dcn.2016.11.006>.
- Eack, S.M., Newhill, C.E., 2012. Racial disparities in mental health outcomes after psychiatric hospital discharge among individuals with severe mental illness. *Soc. Work Res.* 36 (1), 41–52. <https://doi.org/10.1093/swr/svs014>.
- Ellis, B.J., Figueredo, A.J., Brumbach, B.H., Schlomer, G.L., 2009. Fundamental dimensions of environmental risk. *Hum. Nat.* 20 (2), 204–268. <https://doi.org/10.1007/s12110-009-9063-7>.
- Elton, A., Tripathi, S.P., Mletzko, T., Young, J., Cisler, J.M., James, G.A., Kiltz, C.D., 2014. Childhood maltreatment is associated with a sex-dependent functional reorganization of a brain inhibitory control network. *Hum. Brain Mapp.* 35 (4), 1654–1667. <https://doi.org/10.1002/hbm.22280>.
- Elzinga, B.M., Molendijk, M.L., Oude Voshaar, R.C., Bus, B.A.A., Prickaerts, J., Spinhoven, P., Penninx, B.J.W.H., 2011. The impact of childhood abuse and recent stress on serum brain-derived neurotrophic factor and the moderating role of BDNF Val66Met. *Psychopharmacology* 214 (1), 319–328. <https://doi.org/10.1007/s00213-010-1961-1>.
- Erickson, K.I., Voss, M.W., Prakash, R.S., Basak, C., Szabo, A., Chaddock, L., et al., 2011. Exercise training increases size of hippocampus and improves memory. *Proc. Natl. Acad. Sci.* 108 (7), 3017–3022. <https://doi.org/10.1073/pnas.1015950108>.
- Erickson, K.I., Leckie, R.L., Weinstein, A.M., 2014. Physical activity, fitness, and gray matter volume. *Neurobiol. Aging* 35, S20–S28. <https://doi.org/10.1016/j.neurobiolaging.2014.03.034>.
- Erickson, K.I., Hillman, C., Stillman, C.M., Ballard, R.M., Bloodgood, B., Conroy, D.E., et al., 2019. Physical activity, cognition, and brain outcomes: a review of the 2018 physical activity guidelines. *Med. Sci. Sports Exerc.* 51 (6), 1242–1251. <https://doi.org/10.1249/MSS.0000000000001936>.
- Evans, G.W., Li, D., Whipple, S.S., 2013. Cumulative risk and child development. *Psychol. Bull.* 139 (6), 1342–1396. <https://doi.org/10.1037/a0031808>.
- Farrell, C., Doolin, K., O'Leary, N., Jairaj, C., Roddy, D., Tozzi, L., et al., 2018. DNA methylation differences at the glucocorticoid receptor gene in depression are related to functional alterations in hypothalamic-pituitary-adrenal axis activity and to early life emotional abuse. *Psychiatry Res.* 265, 341–348. <https://doi.org/10.1016/j.psychres.2018.04.064>.
- Felitti, V.J., Anda, R.F., Nordenberg, D., Williamson, D.F., Spitz, A.M., Edwards, V., et al., 1998. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the adverse childhood experiences (ACE) study. *Am. J. Prev. Med.* 14 (4), 245–258. [https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8).
- Ferrari, L., Vicenzi, M., Tarantini, L., Barretta, F., Sironi, S., Baccarelli, A.A., et al., 2019. Effects of physical exercise on endothelial function and DNA methylation. *Int. J. Environ. Res. Public Health* 16 (14). <https://doi.org/10.3390/ijerph16142530>.
- Firth, J., Stubbs, B., Vancampfort, D., Schuch, F., Lagopoulos, J., Rosenbaum, S., Ward, P.B., 2018. Effect of aerobic exercise on hippocampal volume in humans: a systematic review and meta-analysis. *NeuroImage* 166, 230–238. <https://doi.org/10.1016/j.neuroimage.2017.11.007>.
- Fong, D.Y.T., Ho, J.W.C., Hui, B.P.H., Lee, A.M., Macfarlane, D.J., Leung, S.S.K., et al., 2012. Physical activity for cancer survivors: meta-analysis of randomised controlled trials. *BMJ* 344. <https://doi.org/10.1136/bmj.e70> (jan30 5), e70–e70.
- Fox, S.E., Levitt, P., Ili, C.A.N., 2010. How the timing and quality of early experiences influence the development of brain architecture. *Child Dev.* 81 (1), 28–40. <https://doi.org/10.1111/j.1467-8624.2009.01380.x>.

- Freitas, P.D., Ferreira, P.G., Silva, A.G., Stelmach, R., Carvalho-Pinto, R.M., Fernandes, F. L.A., et al., 2017. The role of exercise in a weight-loss program on clinical control in obese adults with asthma. A randomized controlled trial. *Am. J. Respir. Crit. Care Med.* 195 (1), 32–42. <https://doi.org/10.1164/rccm.201603-0446OC>.
- Friedman, E.M., Karlamangla, A.S., Gruenewald, T., Koretz, B., Seeman, T.E., 2015. Early life adversity and adult biological risk profiles. *Psychosom. Med.* 77 (2), 176–185. <https://doi.org/10.1097/PSY.0000000000000147>.
- Frodl, T., Carballedo, A., Fagan, A.J., Liseicka, D., Ferguson, Y., Meaney, J.F., 2012. Effects of early-life adversity on white matter diffusivity changes in patients at risk for major depression. *J. Psychiatry Neurosci.* 37 (1), 37–45. <https://doi.org/10.1503/jpn.110028>.
- Fuentes, G.C., Castañer, O., Warnberg, J., Subirana, I., Buil-Cosiales, P., Salas-Salvado, J., et al., 2020. Prospective association of physical activity and inflammatory biomarkers in older adults from the PREDIMED-Plus study with overweight or obesity and metabolic syndrome. *Clin. Nutr.* <https://doi.org/10.1016/j.clnu.2020.01.015>.
- Fuller-Thomson, E., Lewis, D.A., 2015. The relationship between early adversities and attention-deficit/hyperactivity disorder. *Child Abuse Negl.* 47, 94–101. <https://doi.org/10.1016/j.chiabu.2015.03.005>.
- Geoffroy, M.-C., Pinto Pereira, S., Li, L., Power, C., 2016. Child neglect and maltreatment and childhood-to-Adulthood cognition and mental health in a prospective birth cohort. *J. Am. Acad. Child Adolesc. Psychiatry* 55 (1), 33–40. <https://doi.org/10.1016/j.jaac.2015.10.012.e3>.
- Gerber, M., Ludyga, S., Mücke, M., Colledge, F., Brand, S., Pühse, U., 2017. Low vigorous physical activity is associated with increased adrenocortical reactivity to psychosocial stress in students with high stress perceptions. *Psychoneuroendocrinology* 80, 104–113. <https://doi.org/10.1016/j.psyneuen.2017.03.004>.
- Gheorghie, D.A., Li, C., Gallacher, J., Bauermeister, S., 2020. Associations of perceived adverse lifetime experiences with brain structure in UK Biobank participants. *J. Child Psychol. Psychiatry*. <https://doi.org/10.1111/jcpp.13298>.
- Gianaros, P.J., Jennings, J.R., Sheu, L.K., Greer, P.J., Kuller, L.H., Matthews, K.A., 2007. Prospective reports of chronic life stress predict decreased grey matter volume in the hippocampus. *NeuroImage* 35 (2), 795–803. <https://doi.org/10.1016/j.neuroimage.2006.10.045>.
- Gianaros, P.J., Horenstein, J.A., Hariri, A.R., Sheu, L.K., Manuck, S.B., Matthews, K.A., Cohen, S., 2008. Potential neural embedding of parental social standing. *Soc. Cogn. Affect. Neurosci.* 3 (2), 91–96. <https://doi.org/10.1093/scan/nsn003>.
- Gianaros, P.J., Manuck, S.B., Sheu, L.K., Kuan, D.C.H., Votruba-Drzal, E., Craig, A.E., Hariri, A.R., 2011. Parental education predicts corticostriatal functionality in adulthood. *Cereb. Cortex* 21 (4), 896–910. <https://doi.org/10.1093/cercor/bhq160>.
- Gonzalez, A., Jenkins, J.M., Steiner, M., Fleming, A.S., 2009. The relation between early life adversity, cortisol awakening response and diurnal salivary cortisol levels in postpartum women. *Psychoneuroendocrinology* 34 (1), 76–86. <https://doi.org/10.1016/j.psyneuen.2008.08.012>.
- Gorka, A.X., Hanson, J.L., Radtke, S.R., Hariri, A.R., 2014. Reduced hippocampal and medial prefrontal gray matter mediate the association between reported childhood maltreatment and trait anxiety in adulthood and predict sensitivity to future life stress. *Biol. Mood Anxiety Disord.* 4, 12. <https://doi.org/10.1186/2045-5380-4-12>.
- Gothe, N.P., Keswani, R.K., McAuley, E., 2016. Yoga practice improves executive function by attenuating stress levels. *Biol. Psychol.* 121, 109–116. <https://doi.org/10.1016/j.biopsycho.2016.10.010>.
- Graignic-Philippe, R., Dayan, J., Chokron, S., Jacquet, A.-Y., Tordjman, S., 2014. Effects of prenatal stress on fetal and child development: a critical literature review. *Neurosci. Biobehav. Rev.* 43, 137–162. <https://doi.org/10.1016/j.neubiorev.2014.03.022>.
- Greenfield, E.A., Moorman, S.M., 2019. Childhood socioeconomic status and later life cognition: evidence from the Wisconsin longitudinal study. *J. Aging Health* 31 (9), 1589–1615. <https://doi.org/10.1177/0898264318783489>.
- Greicius, M.D., Krasnow, B., Reiss, A.L., Menon, V., 2003. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc. Natl. Acad. Sci.* 100 (1), 253–258. <https://doi.org/10.1073/pnas.0135058100>.
- Greicius, M.D., Srivastava, G., Reiss, A.L., Menon, V., 2004. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc. Natl. Acad. Sci.* 101 (13), 4637–4642. <https://doi.org/10.1073/pnas.0308627101>.
- Greicius, M.D., Flores, B.H., Menon, V., Glover, G.H., Solvason, H.B., Kenna, H., et al., 2007. Resting-state functional connectivity in major depression: abnormally increased contributions from subgenual cingulate cortex and thalamus. *Biol. Psychiatry* 62 (5), 429–437. <https://doi.org/10.1016/j.biopsych.2006.09.020>.
- Groot, C., Hooghiemstra, A.M., Rajmakers, P.G.H.M., van Berckel, B.N.M., Scheltens, P., Scherder, E.J.A., et al., 2016. The effect of physical activity on cognitive function in patients with dementia: a meta-analysis of randomized control trials. *Ageing Res. Rev.* 25, 13–23. <https://doi.org/10.1016/j.arr.2015.11.005>.
- Hagstrom, A.D., Marshall, P.W.M., Lonsdale, C., Papalia, S., Cheema, B.S., Toben, C., et al., 2016. The effect of resistance training on markers of immune function and inflammation in previously sedentary women recovering from breast cancer: a randomized controlled trial. *Breast Cancer Res. Treat.* 155 (3), 471–482. <https://doi.org/10.1007/s10549-016-3688-0>.
- Halonen, J.I., Kivimäki, M., Vahtera, J., Pentti, J., Virtanen, M., Ervasti, J., et al., 2017. Childhood adversity, adult socioeconomic status and risk of work disability: a prospective cohort study. *Occup. Environ. Med.* 74 (9), 659–666. <https://doi.org/10.1136/oemed-2017-104319>.
- Hamer, M., Sabia, S., Batty, G.D., Shipley, M.J., Tabák, A.G., Singh-Manoux, A., Kivimäki, M., 2012. Physical activity and inflammatory markers over 10 years follow up in men and women from the Whitehall II cohort study. *Circulation* 126 (8), 928–933. <https://doi.org/10.1161/CIRCULATIONAHA.112.103879>.
- Hanson, J.L., Knodt, A.R., Brigidi, B.D., Hariri, A.R., 2015. Lower structural integrity of the uncinate fasciculus is associated with a history of child maltreatment and future psychological vulnerability to stress. *Dev. Psychopathol.* 27 (4pt2), 1611–1619. <https://doi.org/10.1017/S0954579415000978>.
- Hanson, J.L., Albert, D., Iselin, A.-M.R., Carré, J.M., Dodge, K.A., Hariri, A.R., 2016. Cumulative stress in childhood is associated with blunted reward-related brain activity in adulthood. *Soc. Cogn. Affect. Neurosci.* 11 (3), 405–412. <https://doi.org/10.1093/scan/nsv124>.
- Hanson, J.L., Knodt, A.R., Brigidi, B.D., Hariri, A.R., 2018. Heightened connectivity between the ventral striatum and medial prefrontal cortex as a biomarker for stress-related psychopathology: understanding interactive effects of early and more recent stress. *Psychol. Med.* 48 (11), 1835–1843. <https://doi.org/10.1017/S0033291717003348>.
- Harms, M.B., Birn, R., Provencal, N., Wiechmann, T., Binder, E.B., Giakas, S.W., et al., 2017. Early life stress, FKBP5 methylation, and inhibition-related prefrontal function: a prospective longitudinal study. *Dev. Psychopathol.* 29 (5), 1895–1903. <https://doi.org/10.1017/S095457941700147X>.
- Hawkins, M., Ciciolla, L., Colaizzi, J., Keirns, N., Smith, C., Stout, M., et al., 2020. Adverse childhood experiences and cognitive function among adults with excess adiposity. *Obes. Sci. Pract.* 6 (1), 47–56. <https://doi.org/10.1002/osp4.385>.
- Healy, C., Eaton, A., Cotter, L., Carter, E., Dhondt, N., Cannon, M., 2021. Mediators of the longitudinal relationship between childhood adversity and late adolescent psychopathology. *Psychol. Med.* 1–9. <https://doi.org/10.1017/S0033291721000477>.
- Heinonen, E., Knekt, P., Härkänen, T., Virtala, E., Lindfors, O., 2018. Associations of early childhood adversities with mental disorders, psychological functioning, and suitability for psychotherapy in adulthood. *Psychiatry Res.* 264, 366–373. <https://doi.org/10.1016/j.psychres.2018.04.011>.
- Heldreth, C.M., Guardino, C.M., Wong, L.H., Schetter, C.D., Shapiro, J.R., Schafer, P., et al., 2016. Childhood racism experiences and postpartum depressive symptoms in african american mothers. *J. Soc. Clin. Psychol.* 35 (10), 840–864. <https://doi.org/10.1521/jscp.2016.35.10.840>.
- Herringa, R.J., Phillips, M.L., Fournier, J.C., Kronhaus, D.M., Germain, A., 2013. Childhood and adult trauma both correlate with dorsal anterior cingulate activation to threat in combat veterans. *Psychol. Med.* 43 (7), 1533–1542. <https://doi.org/10.1017/S0033291712002310>.
- Herzberg, M.P., Gunnar, M.R., 2020. Early life stress and brain function: activity and connectivity associated with processing emotion and reward. *NeuroImage* 209, 116493. <https://doi.org/10.1016/j.neuroimage.2019.116493>.
- Hillman, C.H., Erickson, K.I., Kramer, A.F., 2008. Be smart, exercise your heart: exercise effects on brain and cognition. *Nat. Rev. Neurosci.* 9 (1), 58–65. <https://doi.org/10.1038/nrn2298>.
- Hock, R.S., Bryce, C.P., Fischer, L., First, M.B., Fitzmaurice, G.M., Costa, P.T., Galler, J. R., 2018. Childhood malnutrition and maltreatment are linked with personality disorder symptoms in adulthood: results from a Barbados lifespan cohort. *Psychiatry Res.* 269, 301–308. <https://doi.org/10.1016/j.psychres.2018.05.085>.
- Holz, N.E., Boecker, R., Hohm, E., Zohsel, K., Buchmann, A.F., Blomeyer, D., et al., 2015. The long-term impact of early life poverty on orbitofrontal cortex volume in adulthood: results from a prospective study over 25 years. *Neuropsychopharmacology* 40 (4), 996–1004. <https://doi.org/10.1038/npp.2014.277>.
- Hong, F., Tarullo, A.R., Mercurio, A.E., Liu, S., Cai, Q., Malley-Morrison, K., 2018. Childhood maltreatment and perceived stress in young adults: the role of emotion regulation strategies, self-efficacy, and resilience. *Child Abuse Negl.* 86, 136–146. <https://doi.org/10.1016/j.chiabu.2018.09.014>.
- Jaenisch, R., Bird, A., 2003. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat. Genet.* 33 (3), 245–254. <https://doi.org/10.1038/ng1089>.
- Jankord, R., Herman, J.P., 2008. Limbic regulation of hypothalamo-pituitary-adrenocortical function during acute and chronic stress. *Ann. N. Y. Acad. Sci.* 1148, 64–73. <https://doi.org/10.1196/annals.1410.012>.
- Janusek, L.W., Tell, D., Gaylord-Harden, N., Mathews, H.L., 2017. Relationship of childhood adversity and neighborhood violence to a proinflammatory phenotype in emerging adult African American men: an epigenetic link. *Brain Behav. Immun.* 60, 126–135. <https://doi.org/10.1016/j.bbi.2016.10.006>.
- Jokinen, J., Nordström, P., 2009. HPA axis hyperactivity and cardiovascular mortality in mood disorder inpatients. *J. Affect. Disord.* 116 (1), 88–92. <https://doi.org/10.1016/j.jad.2008.10.025>.
- Jonas, W., Mileva-Seitz, V., Girard, A.W., Biscaglia, R., Kennedy, J.L., Sokolowski, M., et al., 2013. Genetic variation in oxytocin rs2740210 and early adversity associated with postpartum depression and breastfeeding duration. *Genes Brain Behav.* 12 (7), 681–694. <https://doi.org/10.1111/gbb.12069>.
- Jonker, I., Rosmalen, J.G.M., Schoevers, R.A., 2017. Childhood life events, immune activation and the development of mood and anxiety disorders: the TRAILS study. *Transl. Psychiatry* 7 (5). <https://doi.org/10.1038/tp.2017.62> e1112–e1112.
- Joseph, J.J., Golden, S.H., 2017. Cortisol dysregulation: the bidirectional link between stress, depression, and type 2 diabetes mellitus. *Ann. N. Y. Acad. Sci.* 1391 (1), 20–34. <https://doi.org/10.1111/nyas.13217>.
- Karlamangla, A.S., Merkin, S.S., Almeida, D.M., Friedman, E.M., Mogle, J.A., Seeman, T. E., 2019. Early-life adversity and dysregulation of adult diurnal cortisol rhythm. *The Journals of Gerontology: Series B* 74 (1), 160–169. <https://doi.org/10.1093/geronb/gby097>.

- Karoly, L.A., Kilburn, M.R., Cannon, J.S., 2005. Proven Benefits of Early Childhood Interventions. Retrieved from. https://www.rand.org/pubs/research_briefs/RB9145.html.
- Kessler, R.C., McLaughlin, K.A., Green, J.G., Gruber, M.J., Sampson, N.A., Zaslavsky, A.M., et al., 2010. Childhood adversities and adult psychopathology in the WHO world mental health surveys. *Br. J. Psychiatry* 197 (5), 378–385. <https://doi.org/10.1192/bjp.bp.110.080499>.
- Klaperski, S., von Dawans, B., Heinrichs, M., Fuchs, R., 2014. Effects of a 12-week endurance training program on the physiological response to psychosocial stress in men: a randomized controlled trial. *J. Behav. Med.* 37 (6), 1118–1133. <https://doi.org/10.1007/s10865-014-9562-9>.
- Klengel, T., Mehta, D., Anacker, C., Rex-Haffner, M., Pruessner, J.C., Pariante, C.M., et al., 2013. Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat. Neurosci.* 16 (1), 33–41. <https://doi.org/10.1038/nn.3275>.
- Knudsen, E.I., 2004. Sensitive periods in the development of the brain and behavior. *J. Cogn. Neurosci.* 16 (8), 1412–1425. <https://doi.org/10.1162/0898929042304796>.
- Koch, W., Teipel, S., Mueller, S., Benninghoff, J., Wagner, M., Bokke, A.L.W., et al., 2012. Diagnostic power of default mode network resting state fMRI in the detection of Alzheimer's disease. *Neurobiol. Aging* 33 (3), 466–478. <https://doi.org/10.1016/j.neurobiolaging.2010.04.013>.
- Kramer, A.F., Colcombe, S., 2018. Fitness effects on the cognitive function of older adults: a meta-analytic study—revisited. *Perspect. Psychol. Sci.* 13 (2), 213–217. <https://doi.org/10.1177/1745691617707316>.
- Kraynak, T.E., Marsland, A.L., Hanson, J.L., Gianaros, P.J., 2019. Retrospectively reported childhood physical abuse, systemic inflammation, and resting corticolimbic connectivity in midlife adults. *Brain Behav. Immun.* 82, 203–213. <https://doi.org/10.1016/j.bbi.2019.08.186>.
- Kuras, Y.I., Assaf, N., Thoma, M.V., Gianferante, D., Hanlin, L., Chen, X., et al., 2017. Blunted diurnal cortisol activity in healthy adults with childhood adversity. *Front. Hum. Neurosci.* 11 <https://doi.org/10.3389/fnhum.2017.00574>.
- Labonté, B., Suderman, M., Maussion, G., Navaro, L., Yerko, V., Mahar, I., et al., 2012. Genome-wide epigenetic regulation by early-life trauma. *Arch. Gen. Psychiatry* 69 (7), 722–731. <https://doi.org/10.1001/archgenpsychiatry.2011.2287>.
- Lambourne, K., Tomporowski, P., 2010. The effect of exercise-induced arousal on cognitive task performance: a meta-regression analysis. *Brain Res.* 1341, 12–24. <https://doi.org/10.1016/j.brainres.2010.03.091>.
- Lang, J.J., Wolfe Phillips, E., Orpana, H.M., Tremblay, M.S., Ross, R., Ortega, F.B., et al., 2018. Field-based measurement of cardiorespiratory fitness to evaluate physical activity interventions. *Bull. World Health Organ.* 96 (11), 794–796. <https://doi.org/10.2471/BLT.18.213728>.
- Lautenschlager, N.T., Cox, K.L., Flicker, L., Foster, J.K., Bockxmeer, F.M., Xiao, J., et al., 2008. Effect of physical activity on cognitive function in older adults at risk for Alzheimer disease: a randomized trial. *JAMA* 300 (9), 1027–1037. <https://doi.org/10.1001/jama.300.9.1027>.
- Leckie, R.L., Oberlin, L.E., Voss, M.W., Prakash, R.S., Szabo-Reed, A., Chaddock-Heyman, L., et al., 2014. BDNF mediates improvements in executive function following a 1-year exercise intervention. *Front. Hum. Neurosci.* 8 <https://doi.org/10.3389/fnhum.2014.00985>.
- Lee, H.B., Richardson, A.K., Black, B.S., Shore, A.D., Kasper, J.D., Rabins, P.V., 2012. Race and cognitive decline among community-dwelling elders with mild cognitive impairment: findings from the memory and medical care study. *Aging Ment. Health* 16 (3), 372–377. <https://doi.org/10.1080/13607863.2011.609533>.
- Lesuis, S.L., Hooijmakers, L., Korosi, A., de Rooij, S.R., Swaab, D.F., Kessels, H.W., et al., 2018. Vulnerability and resilience to Alzheimer's disease: early life conditions modulate neuropathology and determine cognitive reserve. *Alzheimers Res. Ther.* 10 (1), 95. <https://doi.org/10.1186/s13195-018-0422-7>.
- LeTendre, M.L., Reed, M.B., 2017. The effect of adverse childhood experience on clinical diagnosis of a substance use disorder: results of a nationally representative study. *Subst. Use Misuse* 52 (6), 689–697. <https://doi.org/10.1080/10826084.2016.1253746>.
- Lin, X., Zhang, X., Guo, J., Roberts, C.K., McKenzie, S., Wu, W., et al., 2015. Effects of exercise training on cardiorespiratory fitness and biomarkers of cardiometabolic health: a systematic review and meta-analysis of randomized controlled trials. *J. Am. Heart Assoc.* 4 (7) <https://doi.org/10.1161/JAHA.115.002014>.
- Loprinzi, P.D., 2015. Frequency of moderate-to-vigorous physical activity (MVPA) is a greater predictor of systemic inflammation than total weekly volume of MVPA: implications for physical activity promotion. *Physiol. Behav.* 141, 46–50. <https://doi.org/10.1016/j.physbeh.2015.01.002>.
- Loucks, E.B., Huang, Y.-T., Agha, G., Chu, S., Eaton, C.B., Gilman, S.E., et al., 2016. Epigenetic mediators between childhood socioeconomic disadvantage and mid-life body mass index: the new england family study. *Psychosom. Med.* 78 (9), 1053–1065. <https://doi.org/10.1097/PSY.0000000000000411>.
- Lovaglio, W.R., Farag, N.H., Sorocco, K.H., Acheson, A., Cohoon, A.J., Vincent, A.S., 2013. Early life adversity contributes to impaired cognition and impulsive behavior: studies from the Oklahoma family health patterns project. *Alcohol. Clin. Exp. Res.* 37 (4), 616–623. <https://doi.org/10.1111/acer.12016>.
- Lowe, C.J., Staines, W.R., Hall, P.A., 2017. Effects of moderate exercise on cortical resilience: a transcranial magnetic stimulation study targeting the dorsolateral prefrontal cortex. *Psychosom. Med.* 79 (2), 143–152. <https://doi.org/10.1097/PSY.0000000000000361>.
- Mackes, N.K., Golm, D., Sarkar, S., Kumsta, R., Rutter, M., Fairchild, G., et al., 2020. Early childhood deprivation is associated with alterations in adult brain structure despite subsequent environmental enrichment. *Proc. Natl. Acad. Sci. U.S.A.* 117 (1), 641–649. <https://doi.org/10.1073/pnas.1911264116>.
- Marais, L., Stein, D.J., Daniels, W.M.U., 2009. Exercise increases BDNF levels in the striatum and decreases depressive-like behavior in chronically stressed rats. *Metab. Brain Dis.* 24 (4), 587–597. <https://doi.org/10.1007/s11011-009-9157-2>.
- Marden, J.R., Tchetchen, E.J., Kawachi, I., Glymour, M.M., 2017. Contribution of socioeconomic status at 3 life-course periods to late-life memory function and decline: early and late predictors of dementia risk. *Am. J. Epidemiol.* 186 (7), 805–814. <https://doi.org/10.1093/aje/kwx155>.
- Martinez-Gomez, D., Lavie, C.J., Hamer, M., Cabanas-Sanchez, V., Garcia-Esquinas, E., Pareja-Galeano, H., et al., 2019. Physical activity without weight loss reduces the development of cardiovascular disease risk factors – a prospective cohort study of more than one hundred thousand adults. *Prog. Cardiovasc. Dis.* 62 (6), 522–530. <https://doi.org/10.1016/j.pcad.2019.11.010>.
- Mayat, Z., Giardina, E.-G.V., Liao, M., Aggarwal, B., 2020. Associations of inflammation, physical activity, and sleep in a diverse population of women. *J. Womens Health* 29 (7), 1007–1016. <https://doi.org/10.1089/jwh.2019.7663>.
- McEwen, Bruce S., 2003. Early life influences on life-long patterns of behavior and health. *Ment. Retard. Dev. Disabil. Res. Rev.* 9 (3), 149–154. <https://doi.org/10.1002/mrdd.10074>.
- McEwen, B.S., De Kloet, E.R., Rostene, W., 1986. Adrenal steroid receptors and actions in the nervous system. *Physiol. Rev.* 66 (4), 1121–1188. <https://doi.org/10.1152/physrev.1986.66.4.1121>.
- McGuire, T.G., Miranda, J., 2008. New evidence regarding racial and ethnic disparities in mental health: policy implications. *Health Aff.* 27 (2), 393–403. <https://doi.org/10.1377/hlthaff.27.2.393>.
- McLaughlin, K.A., 2016. Future directions in childhood adversity and youth psychopathology. *J. Clin. Child Adolesc. Psychol.* 45 (3), 361–382. <https://doi.org/10.1080/15374416.2015.1110823>. American Psychological Association, Division 53.
- McLaughlin, K.A., Sheridan, M.A., 2016. Beyond cumulative risk: a dimensional approach to childhood adversity. *Curr. Dir. Psychol. Sci.* 25 (4), 239–245. <https://doi.org/10.1177/0963721416655883>.
- McLaughlin, K.A., Sheridan, M.A., Lambert, H.K., 2014. Childhood adversity and neural development: deprivation and threat as distinct dimensions of early experience. *Neurosci. Biobehav. Rev.* 47, 578–591. <https://doi.org/10.1016/j.neubiorev.2014.10.012>.
- McLaughlin, K.A., Sheridan, M.A., Tibu, F., Fox, N.A., Zeanah, C.H., Nelson, C.A., 2015. Causal effects of the early caregiving environment on development of stress response systems in children. *Proc. Natl. Acad. Sci.* 112 (18), 5637–5642. <https://doi.org/10.1073/pnas.1423363112>.
- McLaughlin, K.A., Koenen, K.C., Bromet, E.J., Karam, E.G., Liu, H., Petukhova, M., et al., 2017. Childhood adversities and post-traumatic stress disorder: evidence for stress sensitization in the World Mental Health Surveys. *Br. J. Psychiatry* 211 (5), 280–288. <https://doi.org/10.1192/bjp.bp.116.197640>.
- McMorris, T., Hale, B.J., 2012. Differential effects of differing intensities of acute exercise on speed and accuracy of cognition: a meta-analytical investigation. *Brain Cogn.* 80 (3), 338–351. <https://doi.org/10.1016/j.bandc.2012.09.001>.
- Mehta, K.M., Yeo, G.W., 2017. Systematic review of dementia prevalence and incidence in United States race/ethnic populations. *Alzheimers Dementia* 13 (1), 72–83. <https://doi.org/10.1016/j.jalz.2016.06.2360>.
- Meijer, O.C., Buurstedde, J.C., Schaaf, M.J.M., 2019. Corticosteroid receptors in the brain: transcriptional mechanisms for specificity and context-dependent effects. *Cell. Mol. Neurobiol.* 39 (4), 539–549. <https://doi.org/10.1007/s10571-018-0625-2>.
- Melrose, R.J., Brewster, P., Marquine, M.J., MacKay-Brandt, A., Reed, B., Farias, S.T., Mungas, D., 2015. Early life development in a multiethnic sample and the relation to late life cognition. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 70 (4), 519–531. <https://doi.org/10.1093/geronb/gbt126>.
- Menezes, A.M.B., Oliveira, P.D., Wehrmeister, F.C., Assunção, M.C.F., Oliveira, I.O., Tovo-Rodrigues, L., et al., 2019. Association of modifiable risk factors and IL-6, CRP, and adiponectin: findings from the 1993 Birth Cohort, Southern Brazil. *PLoS One* 14 (5), e0216202. <https://doi.org/10.1371/journal.pone.0216202>.
- Merrick, M.T., Ford, D.C., Ports, K.A., Guinn, A.S., 2018. Prevalence of adverse childhood experiences from the 2011–2014 behavioral risk factor surveillance system in 23 states. *JAMA Pediatr.* 172 (11), 1038–1044. <https://doi.org/10.1001/jamapediatrics.2018.2537>.
- Merrick, M.T., Ford, D.C., Ports, K.A., Guinn, A.S., Chen, J., Klevens, J., et al., 2019. Vital signs: estimated proportion of adult health problems attributable to adverse childhood experiences and implications for prevention — 25 states, 2015–2017. *Morbidity and Mortality Weekly Report* 68 (44), 999–1005. <https://doi.org/10.15585/mmwr.mm6844e1>.
- Migueles, J.H., Cadenas-Sanchez, C., Esteban-Cornejo, I., Torres-Lopez, L.V., Aadland, E., Chastin, S.F., et al., 2020. Associations of objectively-assessed physical activity and sedentary time with hippocampal gray matter volume in children with Overweight/Obesity. *J. Clin. Med.* 9 (4) <https://doi.org/10.3390/jcm9041080>, 1080.
- Miskolczi, C., Halász, J., Mikics, E., 2019. Changes in neuroplasticity following early-life social adversities: the possible role of brain-derived neurotrophic factor. *Pediatr. Res.* 85 (2), 225–233. <https://doi.org/10.1038/s41390-018-0205-7>.
- Montez, J.K., Hayward, M.D., 2014. Cumulative childhood adversity, educational attainment, and active life expectancy among U.S. Adults. *Demography* 51 (2), 413–435. <https://doi.org/10.1007/s13524-013-0261-x>.
- Mullins, T.S., Campbell, E.M., Hogeveen, J., 2020. Neighborhood deprivation shapes motivational-neurocircuit recruitment in children. *Psychol. Sci.* 31 (7), 881–889. <https://doi.org/10.1177/0956797620929299>.
- Muscattell, K.A., Brosso, S.N., Humphreys, K.L., 2020. Socioeconomic status and inflammation: a meta-analysis. *Mol. Psychiatry* 25 (9), 2189–2199. <https://doi.org/10.1038/s41380-018-0259-2>.

- Mustian, K.M., Janelsins, M.C., Peppone, L.J., Kamen, C.S., Guido, J.J., Heckler, C.E., 2015. EXCAP exercise effects on cognitive impairment and inflammation: a URCC NCORP RCT in 479 cancer patients. *J. Clin. Oncol.* 33 (15 suppl) https://doi.org/10.1200/jco.2015.33.15_suppl.9504, 9504–9504.
- Nagamatsu, L.S., 2012. Resistance training promotes cognitive and functional brain plasticity in seniors with probable mild cognitive impairment. *Arch. Intern. Med.* 172 (8) <https://doi.org/10.1001/archinternmed.2012.379>, 666.
- Nagamatsu, L.S., Chan, A., Davis, J.C., Beattie, B.L., Graf, P., Voss, M.W., et al., 2013. Physical Activity Improves Verbal and Spatial Memory in Older Adults With Probable Mild Cognitive Impairment: a 6-Month Randomized Controlled Trial [Research Article] February 24. <https://doi.org/10.1155/2013/861893>.
- Najar, J., Östling, S., Gudmundsson, P., Sundh, V., Johansson, L., Kern, S., et al., 2019. Cognitive and physical activity and dementia: a 44-year longitudinal population study of women. *Neurology* 92 (12), e1322–e1330. <https://doi.org/10.1212/WNL.0000000000007021>.
- Nakajima, K., Takeoka, M., Mori, M., Hashimoto, S., Sakurai, A., Nose, H., et al., 2010. Exercise effects on methylation of ASC gene. *Int. J. Sports Med.* 31 (09), 671–675. <https://doi.org/10.1055/s-0029-1246140>.
- Nanni, V., Uher, R., Danese, A., 2012. Childhood maltreatment predicts unfavorable course of illness and treatment outcome in depression: a meta-analysis. *Am. J. Psychiatry* 169 (2), 141–151. <https://doi.org/10.1176/appi.ajp.2011.11020335>.
- Nelson, C.A., Zeanah, C.H., Fox, N.A., Marshall, P.J., Smyke, A.T., Guthrie, D., 2007. Cognitive recovery in socially deprived young children: the Bucharest early intervention project. *Science* 318 (5858), 1937–1940. <https://doi.org/10.1126/science.1143921>.
- Nikulina, V., Widom, C.S., 2014. Do race, neglect, and childhood poverty predict physical health in adulthood? A multilevel prospective analysis. *Child Abuse Negl.* 38 (3), 414–424. <https://doi.org/10.1016/j.chiabu.2013.09.007>.
- Nilsson, A., Bergens, O., Kadi, F., 2018. Physical activity alters inflammation in older adults by different intensity levels. *Med. Sci. Sports Exerc.* 50 (7), 1502–1507. <https://doi.org/10.1249/MSS.0000000000001582>.
- Nishida, Y., Hara, M., Higaki, Y., Taguchi, N., Nakamura, K., Nanri, H., et al., 2019. Habitual light-intensity physical activity and ASC methylation in a middle-aged population. *Int. J. Sports Med.* 40 (10), 670–677. <https://doi.org/10.1055/a-0965-1374>.
- Northey, J.M., Rattray, B., Pumpa, K.L., Pryor, D.J., Fraser, M.A., Shaw, M.E., et al., 2020. Objectively measured physical activity is associated with dorsolateral prefrontal cortex volume in older adults. *NeuroImage* 221, 117150. <https://doi.org/10.1016/j.neuroimage.2020.117150>.
- Norton, S., Matthews, F.E., Barnes, D.E., Yaffe, K., Brayne, C., 2014. Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. *Lancet Neurol.* 13 (8), 788–794. [https://doi.org/10.1016/S1474-4422\(14\)70136-X](https://doi.org/10.1016/S1474-4422(14)70136-X).
- Nusslock, R., Miller, G.E., 2016. Early-life adversity and physical and emotional health across the lifespan: a neuroimmune network hypothesis. *Biol. Psychiatry* 80 (1), 23–32. <https://doi.org/10.1016/j.biopsych.2015.05.017>.
- Nyström, M., Neely, G., Hassmén, P., Carlbring, P., 2015. Treating major depression with physical activity: a systematic overview with recommendations. *Cogn. Behav. Ther.* 44, 1–12. <https://doi.org/10.1080/160606073.2015.1015440>.
- Oberlin, L.E., Waiwood, A.M., Cumming, T.B., Marsland, A.L., Bernhardt, J., Erickson, K. I., 2017. Effects of physical activity on poststroke cognitive function. *Stroke* 48 (11), 3093–3100. <https://doi.org/10.1161/STROKEAHA.117.017319>.
- Ozawa, M., Shipley, M., Kivimäki, M., Singh-Manoux, A., Brunner, E.J., 2017. Dietary pattern, inflammation and cognitive decline: the Whitehall II prospective cohort study. *Clin. Nutr.* 36 (2), 506–512. <https://doi.org/10.1016/j.clnu.2016.01.013>.
- Packard, C.J., Bezlyak, V., McLean, J.S., Batty, G.D., Ford, I., Burns, H., et al., 2011. Early life socioeconomic adversity is associated in adult life with chronic inflammation, carotid atherosclerosis, poorer lung function and decreased cognitive performance: a cross-sectional, population-based study. *BMC Public Health* 11 (1). <https://doi.org/10.1186/1471-2458-11-42>, 42.
- Papenberg, G., Ferencz, B., Mangialasche, F., Mecocci, P., Cecchetti, R., Kalpouzos, G., et al., 2016. Physical activity and inflammation: effects on gray-matter volume and cognitive decline in aging. *Hum. Brain Mapp.* 37 (10), 3462–3473. <https://doi.org/10.1002/hbm.23252>.
- Pauly, T., Michalowski, V.I., Nater, U.M., Gerstorf, D., Ashe, M.C., Madden, K.M., Hoppmann, C.A., 2019. Everyday associations between older adults' physical activity, negative affect, and cortisol. *Health Psychol.* 38 (6), 494–501. <https://doi.org/10.1037/hea0000743>. APA PsycArticles® (2212269058; 2019-21771-001).
- Pechtel, P., Pizzagalli, D.A., 2011. Effects of early life stress on cognitive and affective function: an integrated review of human literature. *Psychopharmacology* 214 (1), 55–70. <https://doi.org/10.1007/s00213-010-2009-2>.
- Peters, A.T., Burkhouse, K.L., Kinney, K.L., Phan, K.L., 2019. The roles of early life adversity and rumination in neural response to emotional faces amongst anxious and depressed adults. *Psychol. Med.* 49 (13), 2267–2278. <https://doi.org/10.1017/S0033291718003203>.
- Piercy, K.L., Troiano, R.P., Ballard, R.M., Carlson, S.A., Fulton, J.E., Galuska, D.A., et al., 2018. The physical activity guidelines for americans. *JAMA* 320 (19), 2020–2028. <https://doi.org/10.1001/jama.2018.14854>.
- Pinto Pereira, S.M., Stein Merkin, S., Seeman, T., Power, C., 2019. Understanding associations of early-life adversities with mid-life inflammatory profiles: evidence from the UK and USA. *Brain Behav. Immun.* 78, 143–152. <https://doi.org/10.1016/j.bbi.2019.01.016>.
- Popp, J., Wolfgruber, S., Heuser, I., Peters, O., Hüll, M., Schröder, J., et al., 2015. Cerebrospinal fluid cortisol and clinical disease progression in MCI and dementia of Alzheimer's type. *Neurobiol. Aging* 36 (2), 601–607. <https://doi.org/10.1016/j.neurobiolaging.2014.10.031>.
- Puterman, E., O'Donovan, A., Adler, N.E., Tomiyama, A.J., Kemeny, M., Wolkowitz, O. M., Epel, E., 2011. Physical activity moderates stressor-induced rumination on cortisol reactivity. *Psychosom. Med.* 73 (7), 604–611. <https://doi.org/10.1097/PSY.0b013e318229e1e0>.
- Rabin, J.S., Klein, H., Kim, D.R., Schultz, A.P., Yang, H.-S., Hampton, O., et al., 2019. Associations of physical activity and β -Amyloid with longitudinal cognition and neurodegeneration in clinically normal older adults. *JAMA Neurol.* 76 (10), 1203. <https://doi.org/10.1001/jamaneurol.2019.1879>.
- Racine, N., Zumwalt, K., McDonald, S., Tough, S., Madigan, S., 2020. Perinatal depression: the role of maternal adverse childhood experiences and social support. *J. Affect. Disord.* 263, 576–581. <https://doi.org/10.1016/j.jad.2019.11.030>.
- Real, C.C., Ferreira, A.F.B., Chaves-Kirsten, G.P., Torráo, A.S., Pires, R.S., Britto, L.R.G., 2013. BDNF receptor blockade hinders the beneficial effects of exercise in a rat model of Parkinson's disease. *Neuroscience* 237, 118–129. <https://doi.org/10.1016/j.neuroscience.2013.01.060>.
- Rebar, A.L., Stanton, R., Geard, D., Short, C., Duncan, M.J., Vandelandotte, C., 2015. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychol. Rev.* 9 (3), 366–378. <https://doi.org/10.1080/17437199.2015.1022901>.
- Ridout, K.K., Levandowski, M., Ridout, S.J., Gantz, L., Goonan, K., Palermo, D., et al., 2018. Early life adversity and telomere length: a meta-analysis. *Mol. Psychiatry* 23 (4), 858–871. <https://doi.org/10.1038/mp.2017.26>.
- Rimmele, U., Seiler, R., Marti, B., Wirtz, P.H., Ehlert, U., Heinrichs, M., 2009. The level of physical activity affects adrenal and cardiovascular reactivity to psychosocial stress. *Psychoneuroendocrinology* 34 (2), 190–198. <https://doi.org/10.1016/j.psyneuen.2008.08.023>.
- Rosenbaum, S., Tiedemann, A., Sherrington, C., Curtis, J., Ward, P., 2014. Physical activity interventions for people with mental illness: a systematic review and meta-analysis. *J. Sci. Med. Sport* 18, e150. <https://doi.org/10.1016/j.jsams.2014.11.161>.
- Rovio, S., Spulver, G., Nieminen, L.J., Niskanen, E., Winblad, B., Tuomilehto, J., et al., 2010. The effect of midlife physical activity on structural brain changes in the elderly. *Neurobiol. Aging* 31 (11), 1927–1936. <https://doi.org/10.1016/j.neurobiolaging.2008.10.007>.
- Rueggeberg, R., Wrosch, C., Miller, G.E., 2012. The different roles of perceived stress in the association between older adults' physical activity and physical health. *Health Psychol.* 31 (2), 164–171. <https://doi.org/10.1037/a0025242>.
- Ruparel, N., Chai, J.T., Fisher, E.A., Choudhury, R.P., 2017. Inflammatory processes in cardiovascular disease: a route to targeted therapies. *Nat. Rev. Cardiol.* 14 (3), 133–144. <https://doi.org/10.1038/nrcardio.2016.185>.
- Sabia, S., Dugravot, A., Dartigues, J.-F., Abell, J., Elbaz, A., Kivimäki, M., Singh-Manoux, A., 2017. Physical activity, cognitive decline, and risk of dementia: 28 year follow-up of Whitehall II cohort study. *BMJ* 357. <https://doi.org/10.1136/bmj.j2709>.
- Santalla, D.F., Balardin, J.B., Afonso, R.F., Giorjani, G.M., Sato, J.R., Lacerda, S.S., et al., 2019. Greater anteroposterior default mode network functional connectivity in long-term elderly yoga practitioners. *Front. Aging Neurosci.* 11 <https://doi.org/10.3389/fnagi.2019.00158>.
- Sarubin, N., Nothdurfter, C., Schüle, C., Lieb, M., Uhr, M., Born, C., et al., 2014. The influence of Hatha yoga as an add-on treatment in major depression on hypothalamic-pituitary-adrenal-axis activity: a randomized trial. *J. Psychiatr. Res.* 53, 76–83. <https://doi.org/10.1016/j.jpsychires.2014.02.022>.
- Savla, J.T., Roberto, K.A., Jaramillo-Sierra, A.L., Gambrel, L.E., Karimi, H., Butner, L.M., 2013. Childhood abuse affects emotional closeness with family in mid- and later life. *Child Abuse Negl.* 37 (6), 388–399. <https://doi.org/10.1016/j.chiabu.2012.12.009>.
- Saxton, J.M., Scott, E.J., Daley, A.J., Woodroffe, M., Mutrie, N., Crank, H., et al., 2014. Effects of an exercise and hypocaloric healthy eating intervention on indices of psychological health status, hypothalamic-pituitary-adrenal axis regulation and immune function after early-stage breast cancer: a randomised controlled trial. *Breast Cancer Research: BCR* 16 (2), R39. <https://doi.org/10.1186/bcr3643>.
- Schmidt, M.E., Meynköhn, A., Habermann, N., Wiskemann, J., Oelmann, J., Hof, H., et al., 2016. Resistance exercise and inflammation in breast cancer patients undergoing adjuvant radiation therapy: mediation analysis from a randomized, controlled intervention trial. *Int. J. Radiat. Oncol.* 94 (2), 329–337. <https://doi.org/10.1016/j.ijrobp.2015.10.058>.
- Schuch, F.B., Vancampfort, D., Firth, J., Rosenbaum, S., Ward, P.B., Silva, E.S., et al., 2018. Physical activity and incident depression: a meta-analysis of prospective cohort studies. *Am. J. Psychiatry* 175 (7), 631–648. <https://doi.org/10.1176/appi.ajp.2018.17111194>.
- Seo, D., Rabinowitz, A.G., Douglas, R.J., Sinha, R., 2019. Limbic response to stress linking life trauma and hypothalamus-pituitary-adrenal axis function. *Psychoneuroendocrinology* 99, 38–46. <https://doi.org/10.1016/j.psyneuen.2018.08.023>.
- Shalev, I., Belsky, J., 2016. Early-life stress and reproductive cost: A two-hit developmental model of accelerated aging? *Med. Hypotheses* 90, 41–47. <https://doi.org/10.1016/j.mehy.2016.03.002>.
- Shanmugan, S., Loughead, J., Cao, W., Sarnell, M.D., Satterthwaite, T.D., Ruparel, K., et al., 2017. Impact of tryptophan depletion on executive system function during menopause is moderated by childhood adversity. *Neuropsychopharmacology* 42 (12), 2398–2406. <https://doi.org/10.1038/npp.2017.64>.
- Shanmugan, S., Sarnell, M.D., Loughead, J., Ruparel, K., Gur, R.C., Brown, T.E., et al., 2020. Executive function after risk-reducing salpingo-oophorectomy in BRCA1 and BRCA2 mutation carriers: Does current mood and early life adversity matter? *Menopause* 27 (7), 746–755. <https://doi.org/10.1097/GME.0000000000001535>.
- Shapiro, B.G., Weiss, R.B., Burke, T.A., Boland, E.M., Abramson, L.Y., Alloy, L.B., 2017. Kindling of life stress in bipolar disorder: effects of early adversity. *Behav. Ther.* 48 (3), 322–334. <https://doi.org/10.1016/j.beth.2016.12.003>.

- Sheridan, M.A., McLaughlin, K.A., 2014. Dimensions of early experience and neural development: deprivation and threat. *Trends Cogn. Sci.* 18 (11), 580–585. <https://doi.org/10.1016/j.tics.2014.09.001>.
- Short, A.K., Baram, T.Z., 2019. Early-life adversity and neurological disease: age-old questions and novel answers. *Nat. Rev. Neurol.* 15 (11), 657–669. <https://doi.org/10.1038/s41582-019-0246-5>.
- Sink, K.M., Espeland, M.A., Castro, C.M., Church, T., Cohen, R., Dodson, J.A., et al., 2015. Effect of a 24-Month physical activity intervention vs health education on cognitive outcomes in sedentary older adults: the LIFE randomized trial. *JAMA* 314 (8), 781–790. <https://doi.org/10.1001/jama.2015.9617>.
- Smith, J.C., Nielson, K.A., Woodard, J.L., Seidenberg, M., Durgerian, S., Hazlett, K.E., et al., 2014. Physical activity reduces hippocampal atrophy in elders at genetic risk for Alzheimer's disease. *Front. Aging Neurosci.* 6 <https://doi.org/10.3389/fnagi.2014.00061>.
- Smith, J.A., Zhao, W., Wang, X., Ratliff, S.M., Mukherjee, B., Kardia, S.L.R., et al., 2017. Neighborhood characteristics influence DNA methylation of genes involved in stress response and inflammation: the Multi-Ethnic Study of Atherosclerosis. *Epigenetics* 12 (8), 662–673. <https://doi.org/10.1080/15592294.2017.1341026>.
- Sothmann, M.S., Buckworth, J., Claytor, R.P., Cox, R.H., White-Welkley, J.E., Dishman, R.K., 1996. Exercise training and the cross-stressor adaptation hypothesis. *Exerc. Sport Sci. Rev.* 24 (1). Retrieved from https://journals.lww.com/acsm-essr/Fulltext/1996/00240/Exercise_Training_and_the_Cross_Stressor.11.aspx.
- Spencer, R.L., Deak, T., 2017. A users guide to HPA axis research. *Physiol. Behav.* 178, 43–65. <https://doi.org/10.1016/j.physbeh.2016.11.014>.
- Sripada, R.K., Swain, J.E., Evans, G.W., Welsh, R.C., Liberzon, I., 2014. Childhood poverty and stress reactivity are associated with aberrant functional connectivity in default mode network. *Neuropsychopharmacology* 39 (9), 2244–2251. <https://doi.org/10.1038/npp.2014.75>.
- Stern, Y., Gurland, B., Tatemichi, T.K., Tang, M.X., Wilder, D., Mayeux, R., 1994. Influence of education and occupation on the incidence of alzheimer's disease. *JAMA* 271 (13), 1004–1010. <https://doi.org/10.1001/jama.1994.03510370056032>.
- Stewart, J.C., Rand, K.L., Muldoon, M.F., Kamarck, T.W., 2009. A prospective evaluation of the directionality of the depression–inflammation relationship. *Brain Behav. Immun.* 23 (7), 936–944. <https://doi.org/10.1016/j.bbi.2009.04.011>.
- Stillman, C.M., Lopez, O.L., Becker, J.T., Kuller, L.H., Mehta, P.D., Tracy, R.P., Erickson, K.I., 2017. Physical activity predicts reduced plasma β amyloid in the Cardiovascular Health Study. *Ann. Clin. Transl. Neurol.* 4 (5), 284–291. <https://doi.org/10.1002/acn3.397>.
- Stillman, C.M., Donofry, S.D., Erickson, K.I., 2019. Exercise, fitness and the aging brain: a review of functional connectivity in aging. *Arch. Psychol.* 3 (4) <https://doi.org/10.31296/aop.v3i4.98>.
- Strahler, J., Fuchs, R., Nater, U.M., Klaperski, S., 2016. Impact of physical fitness on salivary stress markers in sedentary to low-active young to middle-aged men. *Psychoneuroendocrinology* 68, 14–19. <https://doi.org/10.1016/j.psyneuen.2016.02.022>.
- Su, S., Jimenez, M.P., Roberts, C.T.F., Loucks, E.B., 2015. The role of adverse childhood experiences in cardiovascular disease risk: a review with emphasis on plausible mechanisms. *Curr. Cardiol. Rep.* 17 (10), 88. <https://doi.org/10.1007/s11886-015-0645-1>.
- Suetani, S., Mamun, A., Williams, G.M., Najman, J.M., McGrath, J.J., Scott, J.G., 2017. Longitudinal association between physical activity engagement during adolescence and mental health outcomes in young adults: a 21-year birth cohort study. *J. Psychiatr. Res.* 94, 116–123. <https://doi.org/10.1016/j.jpsychires.2017.06.013>.
- Sumner, J.A., Colich, N.L., Uddin, M., Armstrong, D., McLaughlin, K.A., 2019. Early experiences of threat, but not deprivation, are associated with accelerated biological aging in children and adolescents. *Biol. Psychiatry* 85 (3), 268–278. <https://doi.org/10.1016/j.biopsych.2018.09.008>.
- Szuhany, K.L., Bugatti, M., Otto, M.W., 2015. A meta-analytic review of the effects of exercise on brain-derived neurotrophic factor. *J. Psychiatr. Res.* 60, 56–64. <https://doi.org/10.1016/j.jpsychires.2014.10.003>.
- Tartar, J., Salzmann, S., Pierreulou, R., Antonio, J., 2018. Acute aerobic exercise decreases a neurophysiological response to emotionally negative stimuli. *OPEN ACCESS* 1 (5), 10.
- Tartibian, B., FitzGerald, L.Z., Azadpour, N., Maleki, B.H., 2015. A randomized controlled study examining the effect of exercise on inflammatory cytokine levels in post-menopausal women. *Post Reprod. Health* 21 (1), 9–15. <https://doi.org/10.1177/2053369114565708>.
- Teicher, M.H., Anderson, C.M., Polcari, A., 2012. Childhood maltreatment is associated with reduced volume in the hippocampal subfields CA3, dentate gyrus, and subiculum. *Proc. Natl. Acad. Sci.* 109 (9), E563–E572. <https://doi.org/10.1073/pnas.1115396109>.
- Thoenen, H., 1995. Neurotrophins and neuronal plasticity. *Science* 270 (5236), 593–598. *JSTOR*. Retrieved from *JSTOR*.
- Thomas, A.G., Dennis, A., Rawlings, N.B., Stagg, C.J., Matthews, L., Morris, M., et al., 2016. Multi-modal characterization of rapid anterior hippocampal volume increase associated with aerobic exercise. *NeuroImage* 131, 162–170. <https://doi.org/10.1016/j.neuroimage.2015.10.090>.
- Tobi, E.W., Sliker, R.C., Luijk, R., Dekkers, K.F., Stein, A.D., Xu, K.M., et al., 2018. DNA methylation as a mediator of the association between prenatal adversity and risk factors for metabolic disease in adulthood. *Sci. Adv.* 4 (1), ea04364 <https://doi.org/10.1126/sciadv.aao4364>.
- Tomeleri, C.M., Ribeiro, A.S., Souza, M.F., Schiavoni, D., Schoenfeld, B.J., Venturini, D., et al., 2016. Resistance training improves inflammatory level, lipid and glycemic profiles in obese older women: a randomized controlled trial. *Exp. Gerontol.* 84, 80–87. <https://doi.org/10.1016/j.exger.2016.09.005>.
- Topel, M.L., Kelli, H.M., Lewis, T.T., Dunbar, S.B., Vaccarino, V., Taylor, H.A., Quyyumi, A.A., 2018. High neighborhood incarceration rate is associated with cardiometabolic disease in nonincarcerated black individuals. *Ann. Epidemiol.* 28 (7), 489–492. <https://doi.org/10.1016/j.annepidem.2018.01.011>.
- Tregellas, J.R., Wylie, K.P., Rojas, D.C., Tanabe, J., Martin, J., Kronberg, E., et al., 2011. Altered default network activity in obesity. *Obesity* 19 (12), 2316–2321. <https://doi.org/10.1038/oby.2011.119>.
- Tyrka, A.R., Price, L.H., Marsit, C., Walters, O.C., Carpenter, L.L., 2012. Childhood adversity and epigenetic modulation of the leukocyte glucocorticoid receptor: preliminary findings in healthy adults. *PLoS One* 7 (1), e30148. <https://doi.org/10.1371/journal.pone.0030148>.
- Underwood, M.D., Bakalian, M.J., Escobar, T., Kassir, S., Mann, J.J., Arango, V., 2019. Early-life adversity, but not suicide, is associated with less prefrontal cortex gray matter in adulthood. *Int. J. Neuropsychopharmacol.* 22 (5), 349–357. <https://doi.org/10.1093/ijnp/pyz013>.
- Unternehmer, E., Meyer, A.H., Burkhardt, S.C.A., Dempster, E., Staehli, S., Theill, N., et al., 2015. Childhood maternal care is associated with DNA methylation of the genes for brain-derived neurotrophic factor (BDNF) and oxytocin receptor (OXTR) in peripheral blood cells in adult men and women. *Stress* 18 (4), 451–461. <https://doi.org/10.3109/10253890.2015.1038992>.
- Vaiserman, A.M., Koliada, A.K., 2017. Early-life adversity and long-term neurobehavioral outcomes: Epigenome as a bridge? *Hum. Genomics* 11 (1), 34. <https://doi.org/10.1186/s40246-017-0129-z>.
- van Bodegom, M., Homberg, J.R., Henckens, M.J.A.G., 2017. Modulation of the hypothalamic-pituitary-Adrenal Axis by early life stress exposure. *Front. Cell. Neurosci.* 11 <https://doi.org/10.3389/fncel.2017.00087>.
- Vaynman, S., Ying, Z., Gomez-Pinilla, F., 2004. Hippocampal BDNF mediates the efficacy of exercise on synaptic plasticity and cognition. *Eur. J. Neurosci.* 20 (10), 2580–2590. <https://doi.org/10.1111/j.1460-9568.2004.03720.x>.
- Vella, C.A., Allison, M.A., Cushman, M., Jenny, N.S., Miles, M.P., Larsen, B., et al., 2017. Physical activity and adiposity-related inflammation: the MESA. *Med. Sci. Sports Exerc.* 49 (5), 915–921. <https://doi.org/10.1249/MSS.0000000000001179>.
- Voss, M.W., Prakash, R.S., Erickson, K.I., Basak, C., Chaddock, L., Kim, J.S., et al., 2010. Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Front. Aging Neurosci.* 2 <https://doi.org/10.3389/fnagi.2010.00032>.
- Voss, M.W., Erickson, K.I., Prakash, R.S., Chaddock, L., Kim, J.S., Alves, H., et al., 2013. Neurobiological markers of exercise-related brain plasticity in older adults. *Brain Behav. Immun.* 28, 90–99. <https://doi.org/10.1016/j.bbi.2012.10.021>.
- Wang, L., Dai, Z., Peng, H., Tan, L., Ding, Y., He, Z., et al., 2014. Overlapping and segregated resting-state functional connectivity in patients with major depressive disorder with and without childhood neglect. *Hum. Brain Mapp.* 35 (4), 1154–1166. <https://doi.org/10.1002/hbm.22241>.
- Wang, D.-C., Lin, H.-T., Lee, Y.-J., Yu, H.-F., Wu, S.-R., Qamar, M.U., 2020. Recovery of BDNF and CB1R in the prefrontal cortex underlying improvement of working memory in prenatal DEHP-Exposed male rats after aerobic exercise. *Int. J. Mol. Sci.* 21 (11) <https://doi.org/10.3390/ijms21113867>, 3867.
- Wanner, B., Vitaro, F., Tremblay, R.E., Turecki, G., 2012. Childhood trajectories of anxiousness and disruptiveness explain the association between early-life adversity and attempted suicide. *Psychol. Med.* 42 (11), 2373–2382. <https://doi.org/10.1017/S0033291712000438>.
- Wassink, J., Perreira, K.M., Harris, K.M., 2017. Beyond Race/Ethnicity: skin color and cardiometabolic health among blacks and hispanics in the United States. *J. Immigr. Minor. Health* 19 (5), 1018–1026. <https://doi.org/10.1007/s10903-016-0495-y>.
- Weinstein, A.M., Voss, M.W., Prakash, R.S., Chaddock, L., Szabo, A., White, S.M., et al., 2012. The association between aerobic fitness and executive function is mediated by prefrontal cortex volume. *Brain Behav. Immun.* 26 (5), 811–819. <https://doi.org/10.1016/j.bbi.2011.11.008>.
- Weuve, J., Barnes, L.L., Mendes de Leon, C.F., Rajan, K.B., Beck, T., Aggarwal, N.T., et al., 2018. Cognitive aging in black and white Americans: cognition, cognitive decline, and incidence of Alzheimer disease dementia. *Epidemiology* 29 (1), 151–159. <https://doi.org/10.1097/EDE.0000000000000747>.
- Whiteman, A.S., Young, D.E., He, X., Chen, T.C., Wagenaar, R.C., Stern, C.E., Schon, K., 2014. Interaction between serum BDNF and aerobic fitness predicts recognition memory in healthy young adults. *Behav. Brain Res.* 259, 302–312. <https://doi.org/10.1016/j.bbr.2013.11.023>.
- Wood, C., Clow, A., Hucklebridge, F., Law, R., Smyth, N., 2017. Physical fitness and prior physical activity are both associated with less cortisol secretion during psychosocial stress. *Anxiety Stress Coping* 31 (2), 135–145.
- Wrann, C.D., White, J.P., Salogiannis, J., Laznik-Bogoslavski, D., Wu, J., Ma, D., et al., 2013. Exercise induces hippocampal BDNF through a PGC-1 α /FNDC5 pathway. *Cell Metab.* 18 (5), 649–659. <https://doi.org/10.1016/j.cmet.2013.09.008>.
- Wunsch, K., Wurst, R., von Dawans, B., Strahler, J., Kasten, N., Fuchs, R., 2019. Habitual and acute exercise effects on salivary biomarkers in response to psychosocial stress. *Psychoneuroendocrinology* 106, 216–225. <https://doi.org/10.1016/j.psyneuen.2019.03.015>.
- Xiong, J., Li, S., Sun, Y., Zhang, X., Dong, Z., Zhong, P., Sun, X., 2015. Long-term treadmill exercise improves spatial memory of male APP^{swe}/PS1^{dE9} mice by regulation of BDNF expression and microglia activation. *Biol. Sport* 32 (4), 295–300. <https://doi.org/10.5604/20831862.1163692>.
- Young-Wolff, K.C., Alabaster, A., McCaw, B., Stoller, N., Watson, C., Sterling, S., et al., 2018. Adverse childhood experiences and mental and behavioral health conditions during pregnancy: the role of resilience. *J. Womens Health* 28 (4), 452–461. <https://doi.org/10.1089/jwh.2018.7108>.
- Yu, Q., Herold, F., Becker, B., KluGah-Brown, B., Zhang, Y., Perrey, S., et al., 2020. Cognitive benefits of exercise interventions: an fMRI activation likelihood estimation

- meta-analysis. *BioRxiv*. <https://doi.org/10.1101/2020.07.04.187401>, 2020.07.04.187401.
- Zalli, A., Jovanova, O., Hoogendijk, W.J.G., Tiemeier, H., Carvalho, L.A., 2016. Low-grade inflammation predicts persistence of depressive symptoms. *Psychopharmacology* 233 (9), 1669–1678. <https://doi.org/10.1007/s00213-015-3919-9>.
- Zeev-Wolf, M., Levy, J., Goldstein, A., Zagoory-Sharon, O., Feldman, R., 2019. Chronic early stress impairs default mode network connectivity in preadolescents and their mothers. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* 4 (1), 72–80. <https://doi.org/10.1016/j.bpsc.2018.09.009>.
- Zeng, H., Irwin, M.L., Lu, L., Risch, H., Mayne, S., Mu, L., et al., 2012. Physical activity and breast cancer survival: an epigenetic link through reduced methylation of a tumor suppressor gene L3MBTL1. *Breast Cancer Res. Treat.* 133 (1), 127–135. <https://doi.org/10.1007/s10549-011-1716-7>.
- Zhang, R., Volkow, N.D., 2019. Brain default-mode network dysfunction in addiction. *NeuroImage* 200, 313–331. <https://doi.org/10.1016/j.neuroimage.2019.06.036>.
- Zhang, F.F., Cardarelli, R., Carroll, J., Zhang, S., Fulda, K.G., Gonzalez, K., et al., 2011. Physical activity and global genomic DNA methylation in a cancer-free population. *Epigenetics* 6 (3), 293–299. <https://doi.org/10.4161/epi.6.3.14378>.
- Zhu, W., Howard, V.J., Wadley, V.G., Hutto, B., Blair, S.N., Vena, J.E., et al., 2015. Association between objectively measured physical activity and cognitive function in older adults—the reasons for geographic and racial differences in stroke study. *J. Am. Geriatr. Soc.* 63 (12), 2447–2454. <https://doi.org/10.1111/jgs.13829>.
- Zschucke, E., Renneberg, B., Dimeo, F., Wüstenberg, T., Ströhle, A., 2015. The stress-buffering effect of acute exercise: evidence for HPA axis negative feedback. *Psychoneuroendocrinology* 51, 414–425. <https://doi.org/10.1016/j.psyneuen.2014.10.019>.