

One Size Fits All? Applying Theoretical Predictions About Age and Emotional Experience to People With Functional Disabilities

Jennifer R. Piazza
California State University, Fullerton

Susan T. Charles
University of California, Irvine

Gloria Luong
Max Planck Institute for Human Development,
Berlin, Germany, and Colorado State University

David M. Almeida
Pennsylvania State University

The current study examined whether commonly observed age differences in affective experience among community samples of healthy adults would generalize to a group of adults who live with significant functional disability. Age differences in daily affect and affective reactivity to daily stressors among a sample of participants with spinal cord injury (SCI) were compared with a noninjured sample. Results revealed that patterns of affective experience varied by sample. Among noninjured adults, older age was associated with lower levels of daily negative affect, higher levels of daily positive affect, and less negative affective reactivity in response to daily stressors. In contrast, among the sample with SCI, no age differences emerged. Findings, which support the model of Strength and Vulnerability Integration, underscore the importance of taking life context into account when predicting age differences in affective well-being.

Keywords: aging, emotion, affect, disability, spinal cord injury

Studies frequently document intact, and even improved, affective well-being across adulthood (see Charles & Carstensen, 2010; Charles & Piazza, 2009, for reviews). The majority of these studies, however, focus predominantly on healthy adults, and it is unclear how physical health problems may alter age-related differences in affective experience. People with physical health problems often face unpredictable, stressful demands due to their condition and are more likely to report affective distress and affective disorders than their healthy counterparts (e.g., Piazza, Charles, & Almeida, 2007; Wolff, Starfield, & Anderson, 2002). Rarely, however, have researchers examined how age-related patterns of affective experience may differ for individuals with health problems, such as functional disabilities. According to the model of Strength and Vulnerability Integration (SAVI), the strength of aging lies in older adults' increased use of strategies that allow

them to avoid highly distressing situations (Charles, 2010). When avoidance of highly stressful situations is difficult or impossible, SAVI predicts that age differences in emotion regulation abilities will be attenuated or disappear completely. The current study tests SAVI using two national samples: a sample with spinal cord injury (SCI) and a sample of adults reporting no functional limitations.

Age and Emotional Experience

A number of studies have documented higher levels of positive affect (PA) and lower levels of negative affect (NA) among older adults compared with younger adults (e.g., Carstensen et al., 2011; Riediger, Schmierek, Wagner, & Lindenberger, 2009; Stone, Schwartz, Broderick, & Deaton, 2010; see Charles & Piazza, 2009, for review). Among the oldest adults, however, age-related patterns sometimes reveal no differences or even upturns in NA and downturns in PA (e.g., Mroczek, 2001). Terminal decline likely accounts for some of these findings (e.g., Gerstorf et al., 2010), but other factors may also be involved.

The SAVI model was formulated to explain how life circumstances influence age differences in emotion regulation strategies and overall levels of affective well-being (Charles, 2010; Charles & Luong, 2013; Charles & Piazza, 2009). According to SAVI, older adults are motivated to maintain affective well-being because of their perception of time left in life (as posited and described by Socioemotional Selectivity Theory; Carstensen, 2006), and are better equipped to do so because of the knowledge and experience acquired from years already lived. Older adults are therefore more likely to avoid exposure to negative experiences (e.g., Blanchard-Fields, 2007) or, when avoidance is impossible, to effectively modify their affective responses to caustic experiences.

This article was published Online First August 31, 2015.

Jennifer R. Piazza, Department of Health Science, California State University, Fullerton; Susan T. Charles, Department of Psychology and Social Behavior, University of California, Irvine; Gloria Luong, Max Planck Research Group "Affect Across the Lifespan," Max Planck Institute for Human Development, Berlin, Germany, and Department of Human Development and Family Studies, Colorado State University; and David M. Almeida, Department of Human Development and Family Studies, Pennsylvania State University.

Supported by the National Institute of Aging and the National Institutes of Health (Grant Nos. R01AG042431, P01AG020166, and R01AG019239).

Correspondence concerning this article should be addressed to Jennifer R. Piazza, 243 KHS, Department of Health Science, California State University, Fullerton, Fullerton, CA 92831. E-mail: jpiazza@fullerton.edu

Emotions in Context

By integrating environmental context and other life circumstances with age-related motivations and abilities, researchers can better predict when age-related improvements in well-being will be most pronounced, and when they will not be present or even reverse in direction. Specifically, some life experiences make it particularly difficult to avoid stressors and/or to modify one's appraisals of stressful events. For example, life experiences such as loss of social belonging, exposure to chronic, uncontrollable stressors, and neurological dysregulation are all associated with high levels of distress. In addition, these experiences (e.g., neurological dysregulation in the form of cognitive impairment) increase in prevalence with age. Thus, although the motivation to maintain affective well-being exists, certain circumstances impinge on older adults' ability to do so (Charles, 2010). SAVI posits, then, that the reason why some studies fail to show age-related improvements in affective well-being is that certain life contexts may prevent older adults from avoiding highly arousing, unpleasant events.

Of course, even though these situations increase in frequency with age (see Hawkey & Cacioppo, 2007; Park, 2002, for reviews), they are not confined to old age. Younger adults can and do experience adverse circumstances such as loss of social belonging (e.g., Leary, 1990; Stroebe, Schut, & Stroebe, 2007) and neurological dysregulation (Salthouse, 2004). What happens, then, to age differences in affective experience among people living in similar difficult circumstances? One study found that older adults with multiple chronic health conditions were just as emotionally reactive to daily stressors as were younger adults who reported the same number of health conditions (Piazza et al., 2007). Although this study lends some support to SAVI, one issue with studying age differences in the context of chronic health conditions is that chronic health conditions are more normative in later adulthood, and thus more of an "on-time" event.

On-time or off-time events refer to the normative timing of positive or negative events at particular stages in life (Neugarten, 1979). Researchers have posited that coping with a negative life event is particularly difficult when it occurs earlier than what is commonly predicted. Indeed, studies have suggested that younger individuals have greater problems adjusting to spousal bereavement (e.g., Stroebe et al., 2007) and chronic illness (Aldwin, Sutton, Chiara, & Spiro, 1996) than do older adults. When an event happens at a more normative age, people may not only have anticipated the event, but may also have same-aged peers living with the same difficult situation. Thus, the influence of the on-time versus off-time phenomena represents confounds in previous investigations of age differences in emotional experience among people faced with problems more common in later life.

The Context of SCI

One life circumstance that is not age-normative—and therefore not subject to the issue of on-time/off-time events—is the occurrence of a spinal cord injury. At no age is living with SCI normative: older adults will not have foreseen, planned, or expected to be living with SCI any more than younger adults. SCI is a severe, disabling condition that affects approximately 273,000 people in the United States, with 12,000 new cases per year (National Spinal Cord Injury Statistical Center, 2013). Living with SCI presents a number of physical and emotional challenges, from

adjusting one's activities due to health and functional constraints to facing potential societal stigma. Rates of emotional disorders among people with SCI are twice that of noninjured adults (Migliorini, Tonge, & Taleporos, 2008) and rates of suicide significantly higher (see Giannini et al., 2010, for a review). In contrast, levels of subjective well-being (Fuhrer, 1996) and life satisfaction are significantly lower (Kemp & Krause, 1999) in this population. Few studies, however, have examined age differences in affective experience among people with SCI.

Affective Reactivity

We were interested in applying SAVI to examine age-related patterns in daily affective experiences, and particularly age differences in affective reactivity in response to daily stressors, among people with SCI compared with people without SCI. Daily stressors, such as an impending work deadline, an argument with a significant other, or a missed appointment, are the hassles that disrupt daily life (Almeida, 2005). Previous findings examining age differences in affective reactivity to daily stressors have been mixed, with some research finding age-related improvements in affective reactivity (e.g., Almeida & Horn, 2004; Birditt, Fingerman, & Almeida, 2005), some reporting null effects (e.g., Diehl & Hay, 2010; Neupert, Almeida, & Charles, 2007), and some finding greater reactivity with age (e.g., Aldwin, Jeong, Igarashi, Choun, & Spiro, 2014; Sliwinski, Almeida, Smyth, & Stawski, 2009). The discrepancies across studies in patterns of age differences in affective reactivity to stressors may be partially explained by the ability of people to disengage from these situations (e.g., Birditt & Fingerman, 2005; Luong & Charles, 2014). In one study, for example, older adults reported less affective distress than younger adults on days when they actively avoided an argument (Charles, Piazza, Luong, & Almeida, 2009). On days when an argument occurred, however, no age-related differences in affective experience emerged. Moreover, younger adults reported equally strong reactivity for both situations, whereas older adults reported less reactivity on days they actively avoided an argument than on days when they reported having an argument.

Avoiding stressful situations may reflect older adults' motivation to structure their daily life in a way that allows for the maintenance of affective well-being (e.g., Blanchard-Fields, 2007; Luong, Charles, & Fingerman, 2011). Indeed, when older adults are unable to use disengagement strategies effectively, such as when they continue to be mentally preoccupied by recent daily hassles, they exhibit increased levels of NA relative to younger individuals (Wrzus, Luong, Wagner, & Riediger, 2015). Research has indicated that people with SCI face not only the same stressors as their noninjured peers but also a number of severe disability-related stressors (Degraff & Schaffer, 2008; Schopp et al., 2007). Moreover, the nature of stressors faced by people with SCI (e.g., bowel/bladder problems, inaccessibility in public places, medical complications) makes avoidance and disengagement from aversive situations very difficult—if not impossible (Degraff & Schaffer, 2008). For example, people with SCI face a number of environmental barriers in their day-to-day life (Lysack, Komanecky, Kabel, Cross, & Neufeld, 2007), including perceived barriers to employment, such as transportation and workplace accessibility (Schopp et al., 2007). They also report difficulty fulfilling social role obligations and completing activities of daily life without

assistance (Noreau & Fougereyrollas, 2000). Thus, for people with SCI, the motivation to avoid or actively disengage from stressors may exist, but the barriers they encounter in their daily life may make it more difficult to do so compared with their same-aged, noninjured peers.

The Current Study

In the current study, we examined age differences in daily affect and affective reactivity to daily stressors among people living with SCI, and compared this pattern with a sample of participants from the National Study of Daily Experiences (NSDE II) who did not have SCI and who reported having no functional limitations. Based on SAVI, we hypothesized that age-related benefits in daily affect and affective reactivity to daily stressors would be present in the noninjured sample (NSDE II), but would be absent or attenuated in the SCI sample. To focus on age differences in relatively normative, daily emotion regulation processes as opposed to age differences in response to adjustment to injury (Krause & Crewe, 1991), we recruited participants with SCI who had been injured for at least 5 years. Moreover, although people with SCI have a life expectancy of about 70–92% of the population life expectancy (Yeo et al., 1998), their condition does not suggest a terminal diagnosis (Garshick et al., 2005). Thus, concerns of terminal decline are reduced for this condition compared with others that have a more life-limiting prognosis.

Method

Participants and Procedure

The NSDE II sample. The NSDE II, which is the daily diary telephone interview portion of the second wave of the Midlife Development in the United States Survey (MIDUS II), was carried out at the same time as the data collection for the SCI sample (described next). NSDE II participants included a national sample of 1,079 MIDUS II participants who were recruited through random-digit phone dialing and had no other family members in the study. Across eight consecutive evenings, NSDE II participants answered questions pertaining to the previous 24 hr, including their physical health, affective well-being, and any stressors they may have encountered. Because the goal of the current study was to examine differences between people with and without functional limitations, analyses were limited to those who reported no problems with activities of daily living (ADLs; described in the Measures section below; $n = 704$). Interviews lasted approximately 20 min each and the first interview was staggered across day of the week. Participants were compensated \$25.00.

The sample with SCI. The SCI sample included 239 participants who were recruited through the Reeve–Irvine Research Center and other SCI-related organizations. Participants on the Reeve–Irvine email listserv were emailed a description of the study with laboratory contact information. Participants were also recruited through SCI email listservs and websites, advertisements placed in regional and national SCI magazines and newsletters, and flyers. To be eligible to participate, individuals were required to be at least 18 years of age, to be free from any psychological disorder, and to have had an SCI for more than 5 years.

On average, participants with SCI had been injured for 20.3 years ($SD = 11.66$) at the time of the initial interview. Level of injury varied across participants: 52% were classified as tetraplegic, having sustained an injury at vertebrate level C7 or higher; 47% were classified as paraplegic, having sustained an injury at vertebrate level T1 or lower; the remaining 1% were unsure of their injury level. Most participants were injured in motor vehicle crashes (40.3%) or other types of accidents (35.4%). Medical problems, such as transverse myelitis and spina bifida, comprised 6.6% of all injuries, and 11.6% of participants were injured in other ways, such as gunshot wounds. Because cause of injury was not assessed in the interview protocol until after the study had already begun, data are missing for 6% of participants. Between 2006 and 2007, participants with SCI completed a series of daily diary telephone interviews, modeled after the protocol used in NSDE II. Prior to the daily diary interviews, participants with SCI completed an initial telephone interview lasting approximately 1 hr, which assessed demographic characteristics and overall physical and affective well-being. Subsequently, they completed a series of daily interviews, during which participants with SCI were asked the same questions as in the NSDE II about their daily physical health, affective well-being, and whether they had experienced any stressors during the previous 24 hr (see Measures section). Due to fiscal constraints, we were unable to interview an SCI sample equivalent in size to the NSDE II sample. To more closely match the samples for total number of days interviewed, we extended SCI interviews to 15 consecutive days prior to the study's inception. This provided sufficient power at the daily level to examine differences between the two samples. After completing the study, participants were compensated \$150.00. Because of the slightly different age ranges of the NSDE II (34–84 years) and SCI samples (21–87 years), analyses for the current study were limited to SCI participants 34 years and older, bringing the final SCI sample size to 181.

Creating a comparison subsample. The NSDE II and SCI samples differed on important demographic variables, so a comparison NSDE II subsample was created. Because of the difference in interview days (15 days for participants with SCI and 8 days for NSDE II participants), two NSDE II participants were retained ($n = 362$) for every participant with SCI ($n = 181$). This ensured that each sample had approximately the same number of interview days for all analyses. Only NSDE II participants who reported no functional limitations were included in this comparison group. Participants were matched according to age, gender, education, and, when possible, ethnicity. Because of the small number of minority participants in NSDE II, however, we could not entirely match on ethnicity.

Average age of participants did not differ between the two samples, $t(541) = .56, p = .52$ (NSDE II: $M_{\text{age}} = 50.7$ years, $SD = 11.0$; SCI: $M_{\text{age}} = 50.3$ years, $SD = 11.5$), and age ranges were nearly identical (NSDE II: 50 years; SCI: 53 years). Both samples had a similar number of females (NSDE: 50.6%; SCI: 54.1), $\chi^2(1, N = 543) = .62, p = .43$, and reported similar levels of education, $t(540) = -.56, p = .58$, with nearly half of participants in both samples reporting having a bachelor's degree or higher (NSDE: 50.3%; SCI: 47.2%). Participants did, however, differ significantly in terms of marital status, $\chi^2(4, N = 541) = 42.9, p < .001$, and ethnicity, $\chi^2(4, N = 543) = 19.6, p < .001$.

For a comparison of demographic characteristics for the NSDE II and SCI samples, see Table 1.

Measures

Activities of daily living. In MIDUS and NSDE II, problems with ADLs were assessed by asking participants to rate on a 4-point scale, ranging from *not at all* to *a lot*, how much their health limited their ability to do each of the following activities: bathing/dressing oneself; climbing one flight of stairs; and walking one block. Of the 1,079 NSDE II participants, 375 reported having some difficulty in one of these domains and were subsequently excluded as possible comparison participants.

Time since injury. Participants in the SCI sample were asked how long they had been injured. Answers ranged from a minimum of 5 years to a maximum of 65 years, with participants injured, on average, for 20.3 years ($SD = 11.66$). Research has indicated that time since injury is associated with greater life adjustment among people with SCI (Krause, 1998). Because age was correlated with time since injury in the current study ($r = .24, p = .01$), time since injury was included as a covariate in all models testing our hypotheses. To ensure that all people were included in the analyses, NSDE II participants were assigned a 0 value for this variable.

Chronic physical health conditions. SCI and NSDE II participants were asked if they had been treated for or experienced 23 chronic health conditions in the past year (e.g., asthma, bronchitis or emphysema, gall bladder trouble, diabetes or high blood sugar; Marmot, Ryff, Bumpass, Shipley, & Marks, 1997), which were subsumed into 17 chronic condition categories. Participants with SCI reported significantly more chronic health conditions than did NSDE II participants, $t(281.2) = -11.2, p < .001$ (NSDE II: 1.38 conditions, $SD = 1.46$; SCI: 3.20 conditions, $SD = 1.97$). In both samples, increasing age was associated with more chronic health conditions (NSDE II: $r = .25, p < .001$; SCI: $r = .29, p < .001$).

Table 1
Sample Characteristics of the NSDE II and SCI Samples

Variables	NSDE II sample (%)	SCI sample (%)
Gender		
Female	50.6	54.1
Education		
< High school	1.9	2.8
High school degree	13.0	12.2
Some college	34.8	37.8
Bachelor's degree	50.3	47.2
Ethnicity*		
White	90.1	76.8
Black	2.8	9.4
Asian	0.8	2.2
Hispanic	3.6	7.7
Other	2.8	3.8
Marital status*		
Married	75.4	48.3
Separated	1.7	2.8
Divorced	10.8	16.7
Widowed	2.5	5.6
Never married	9.7	26.7

Note. NSDE = National Study of Daily Experiences; SCI = spinal cord injury.

* $p < .001$.

For a complete listing of sample differences in chronic conditions, see Table 2.

Daily NA. In both the NSDE II and SCI samples, daily NA was assessed using combined items from the Positive and Negative Affect Schedule (Watson, Clark, & Tellegen, 1988) and the revised Non-Specific Psychological Distress Scale (Kessler et al., 2002). Respondents indicated on a 5-point scale, anchored at 0 (*none of the time*) and 4 (*all of the time*), how much of the time they experienced each of the following emotions or emotion descriptors on each interview day: so sad nothing could cheer you up, nervous, restless/fidgety, hopeless, that everything was an effort, worthless, lonely, afraid, angry, and frustrated. Mean scores for these 10 items were calculated for each participant (for the NSDE II sample, daily α range: .66–.79; for the SCI sample, daily α range: .73–.83).

Daily PA. To assess daily PA in the NSDE II and SCI samples, participants rated on a 5-point scale, anchored at 0 (*none of the time*) and 4 (*all of the time*), how much of the time they experienced each of the following 10 emotions or emotion descriptors on each interview day: cheerful, in good spirits, extremely happy, calm and peaceful, satisfied, full of life, close to others, like you belong, enthusiastic, and proud. Mean scores for these 10 emotions were calculated for each participant, resulting in an average PA score (for the NSDE II sample, daily α range: .91–.94; for the SCI sample, daily α range: .91–.95).

Daily stressors and affective reactivity. For both samples, daily stressors were assessed using the semi-structured Daily Inventory of Stressful Experiences (Almeida, Wethington, & Kessler, 2002). The inventory consists of a series of seven stem questions asking whether certain types of daily stressors, such as a problem in the home or an argument with someone, had occurred in the previous 24 hr. For each daily interview, individuals responding affirmatively to any of the seven stressors were classified as having had experienced a stressor (1) and those who reported not experiencing a stressor were classified as having had experienced no stressors (0).

Person-mean average number of stressors was calculated by averaging the number of stressors each participant reported across the interview period and was used as a covariate in analyses examining reactivity to daily stressors (L. Hoffman & Stawski, 2009).

Affective reactivity to daily stressors was operationalized as the within-person slope between stressors and affect and refers to an individual's change in NA or PA from a nonstressor day to a stressor day.

Results

The first set of analyses examined sample and age differences in average levels of daily NA and PA. The second set of analyses tested sample and age differences in negative and positive affective reactivity to daily stressors. For all analyses, we used two-level hierarchical multilevel modeling (MLM) to examine between- and within-person differences (Raudenbush & Bryk, 2002). In MLM, Level 1 represents within-person variability and Level 2 represents between-person variability. In all models, the Level 1 outcome variable, $affect_{i,t}$, refers to within-person variation in affect (either NA or PA) for person i on day t , and is a function of a person-specific intercept, within-person error, and—for the

Table 2
Percentage of NSDE and SCI Participants Reporting Chronic Health Conditions

Type of chronic condition	NSDE participants % of sample	SCI participants % of sample
Autoimmune disorders	1.1%	1.1%
Bone-related conditions ^a	26.8%	47.0%
Cardiovascular conditions ^b	19.1%	16.6%
Diabetes or high blood sugar	6.1%	12.7%
Digestive conditions ^c	16.9%	49.7%
Foot trouble	8.0%	24.3%
Gall bladder trouble	0.8%	2.8%
Hay fever	11.9%	14.4%
Hernia	1.1%	4.4%
HIV/AIDS	0.0%	0.0%
Lung conditions ^d	7.2%	9.4%
Migraine headaches	6.6%	11.1%
Neurological conditions	1.9%	7.8%
Skin trouble	8.6%	34.3%
Thyroid disease	7.5%	9.9%
Trouble with gums, mouth, or teeth ^e	6.1%	13.8%
Urinary or bladder problems	8.0%	60.8%
% reporting 0 chronic conditions	36.5%	5.5%
% reporting 1 chronic condition	25.7%	16.6%
% reporting multiple chronic conditions	37.8%	77.9%

Note. NSDE = National Study of Daily Experiences; SCI = spinal cord injury.

^a Bone-related conditions include arthritis, rheumatism, or other bone/joint diseases; sciatica, lumbago, or recurring backache. ^b Cardiovascular conditions include stroke, high blood pressure, or hypertension. ^c Digestive conditions include recurring stomach trouble, indigestion, or diarrhea; constipated all/most of time. ^d Lung conditions include asthma, bronchitis, emphysema, tuberculosis, and other lung problems. ^e Trouble with gums, mouth, or teeth include persistent trouble with gums or mouth and persistent trouble with teeth.

stressor reactivity models—stressor exposure, which was included as a random effect.

In all models, Level 2 between-person variables included the sample (SCI or NSDE II), participant’s age, and other demographic variables that differed by sample or that could influence age differences in affective experience: education, race (effect coded with White as the reference group), marital status (effect coded with married as the reference group), gender (effect coded with female as the reference group), interview day (1–8 for NSDE II; 1–15 for SCI), time since injury (NSDE participants were assigned 0 for this variables so that they were not omitted from analyses), and random intraindividual variation. In follow-up analyses examining within-group differences, we also statistically adjusted for injury level in the SCI group. In the stressor reactivity models, we also adjusted for the average number of daily stressors participants reported across the interview period. In subsequent models, interaction terms were included to test the hypotheses. For the daily affect models, we included a Sample × Age interaction term as a test of our hypotheses. For the stressor reactivity models, we included the following interactions: Sample × Age, Stressor × Age, Stressor × Sample, and a three-way Sample × Age × Stressor interaction. In these models, the three-way interaction tested our main hypothesis. Analyses were conducted using SAS PROC MIXED, version 9.3, and estimated from unstructured covariance matrices by means of full maximum likelihood.

Daily NA

Results from analyses examining daily NA are presented in Table 3, Model 1. This model revealed a significant Age × Sample interaction, which indicated that older age was related to lower daily NA in the NSDE II sample, but was unrelated to daily NA in the SCI sample (see Figure 1). To examine this interaction further, we conducted two separate MLMs with daily NA as the outcome variable; the first model examined only the NSDE II participants; the second model examined only participants with SCI. Whereas the NSDE II analyses showed a significant, negative effect of age on NA ($\beta = -.004, p < .001$), the SCI analyses did not ($\beta = .002, p = .26$), even when adjusting for level of injury ($\beta = -.001, p = .55$). Additional findings from the full model indicated that people who had never been married reported higher levels of NA than those who were married ($\beta = .060, p < .05$) and, as the study progressed, people reported less NA (see Table 3, Model 1).

Daily PA

Results from analyses examining daily PA are presented in Table 4, Model 1. In this model, an Age × Sample interaction emerged, which revealed that although older age was related to

Table 3
Age and Sample Differences in Daily Negative Affect and Negative Affective Reactivity

Fixed effects	Model 1 Estimate (SE)	Model 2 Estimate (SE)
Intercept	.197 (.047)***	.056 (.037)
Age	-.004 (.001)***	-.001 (.001)
Sex (ref = female)	-.034 (.020)	.001 (.016)
Sample (ref = NSDE)	.191 (.038)***	.193 (.029)***
Ethnicity (ref = White)		
African American	.006 (.058)	-.000 (.046)
Asian	-.024 (.048)	.027 (.037)
Hispanic	.052 (.091)	.056 (.070)
Other	-.069 (.046)	-.060 (.036)
Education	.011 (.013)	-.005 (.010)
Marital status (ref = married)		
Separated	-.056 (.071)	-.014 (.055)
Divorced	-.008 (.031)	-.019 (.024)
Widowed	.010 (.057)	.017 (.044)
Never married	.060 (.030)*	.055 (.024)*
Day	-.007 (.001)***	-.003 (.001)***
Time since injury	-.003 (.002)	-.004 (.001)***
Age × Sample	.006 (.002)**	.003 (.001)**
Stressor		.151 (.012)***
Average no. of stressors		1.19 (.139)***
Age × Stressor		-.003 (.001)**
Stressors × Sample		.009 (.019)
Stressors × Sample × Age		.003 (.002)*
Random effects		
Intercept	.046 (.003)***	.022 (.002)***
Stressor		.015 (.003)***
Covariance		.010 (.002)***
Residual	.057 (.001)***	.049 (.001)***
Model fit		
AIC	1,053.8	335.2
BIC	1,131.1	442.5

Note. AIC = Akaike information criterion; BIC = Bayesian information criteria; NSDE = National Study of Daily Experiences. * $p < .05$. ** $p < .01$. *** $p < .001$.




Figure 1. Age and sample differences in daily negative affect. SCI = spinal cord injury; NSDE = National Study of Daily Experiences.

higher levels of daily PA in the NSDE II sample, no age differences emerged in the SCI sample (see Figure 2). We once again conducted separate MLMs for each sample to examine this interaction further. Results revealed that increasing age was associated with higher levels of PA in the NSDE sample ($\beta = .015, p < .001$), but not the SCI sample ($\beta = .001, p = .90$), even when adjusting for level of injury ($\beta = .007, p = .28$). Additional findings in the full model revealed that higher levels of education were associated with lower levels of daily PA and that racial groups self-defined as “other” (i.e., not African American, Asian, or Hispanic) reported higher levels of daily PA than did Whites.

Affective Reactivity to Daily Stressors

Two sets of analyses were conducted to examine age and sample differences in affective reactivity in response to daily stressors. The first set of analyses examined negative affective reactivity; the second set of analyses examined positive affective reactivity.

Negative affective reactivity. Results examining negative affective reactivity are presented in Table 3, Model 2. Results revealed a significant three-way Age \times Sample \times Stressor interaction ($\beta = .003, p = .04$; see Figure 3), which indicated that although older age was related to lower negative affective reactivity (i.e., change in NA from a stressor to a nonstressor day) in




Figure 2. Age and sample differences in daily positive affect. SCI = spinal cord injury; NSDE = National Study of Daily Experiences.

the NSDE II sample, it was unrelated to negative affective reactivity in the SCI sample. To examine this interaction further, we conducted additional MLMs: the first examined only the NSDE II participants and the second examined only the participants with SCI. In analyses examining the NSDE II participants, an Age \times Stressor interaction emerged, indicating that, in the NSDE sample, increasing age was related to less negative affective reactivity to daily stressors ($\beta = -.003, p = .003$). In the sample with SCI, however, age was not significantly related to negative affective reactivity ($\beta = .000, p = .72$), even when adjusting for level of injury ($\beta = .001, p = .57$). Findings from the full model also revealed that people who were never married reported higher levels of NA compared with married individuals ($\beta = .055, p = .02$), as did people who reported a greater number of stressors across the week ($\beta = 1.19, p < .001$). In addition, time since injury was inversely associated with levels of NA ($\beta = -.004, p < .001$) and people reported less NA as the interview days progressed.

PA reactivity. No significant interactions emerged in an analysis examining age and sample differences in positive affective reactivity in response to stressors (i.e., change in PA from a stressor day to a nonstressor day; see Table 4, Model 2). However, people who reported more stressors on average (regardless of age and sample) reported lower levels of PA than people reporting fewer stressors. People also reported lower levels of PA on days they encountered stressors.

Discussion

Researchers have offered various explanations for the paradox of well-being in later life, attributing it to reasons such as physiological changes in the brain (Cacioppo, Berntson, Bechara, Tranel, & Hawkey, 2011) and developmental shifts in cognitive processing of information (Labouvie-Vief, 2003; see Isaacowitz & Blanchard-Fields, 2012, for a discussion). The current study approached this question from a contextual perspective, examining how the circumstance of living with a functional disability may alter patterns of age differences in daily affective experiences. Results indicated that, among people with no significant functional




Figure 3. Age and sample differences in negative affective reactivity to daily stressors. SCI = spinal cord injury; NSDE = National Study of Daily Experiences.

Table 4
Age and Sample Differences in Daily Positive Affect and Positive Affective Stressor Reactivity

Fixed effects	Model 1 Estimate (SE)	Model 2 Estimate (SE)
Intercept	2.995 (0.143)***	3.130 (0.140)***
Age	.016 (.003)**	.012 (.004)***
Sex (ref = female)	-.008 (.062)	-.040 (.061)
Sample (ref = NSDE)	-.274 (.117)*	-.277 (.114)*
Ethnicity (ref = White)		
African American	-.119 (.176)	-.074 (.171)
Asian	-.003 (.146)	-.035 (.142)
Hispanic	.099 (.275)	.080 (.267)
Other	.351 (.141)*	.347 (.137)*
Education	-.104 (.040)*	-.067 (.039)
Marital status (ref = married)		
Separated	-.192 (.218)	-.196 (.212)
Divorced	.147 (.095)	.142 (.093)
Widowed	-.029 (.175)	-.046 (.170)
Never married	-.128 (.093)	-.142 (.090)
Day	-.003 (.002)	-.006 (.002)**
Time since injury	.005 (.005)	.005 (.005)
Age × Sample	-.017 (.005)**	-.014 (.006)*
Stressor		-.182 (.022)***
Average no. of stressors		-1.830 (0.464)***
Age × Stressor		.001 (.002)
Stressors × Sample		.044 (.033)
Stressors × Sample × Age		-.002 (.003)
Random effects		
Intercept	.470 (.030)***	.455 (.030)***
Stressor		.025 (.008)***
Covariance		-.022 (.012)
Residual	.198 (.004)***	.188 (.004)***
Model fit		
AIC	8,007.9	7,847.4
BIC	8,085.2	7,954.7

Note. AIC = Akaike information criterion; BIC = Bayesian information criteria; NSDE = National Study of Daily Experiences.

* $p < .05$. ** $p < .01$. *** $p < .001$.

disability, age was associated with higher levels of daily PA, lower levels of daily NA, and less negative affective reactivity in response to daily stressors. Among adults with functional disabilities as a result of SCI, however, no age differences emerged.

Well-Being Among People With a Disability

The physical and functional challenges for people living with a disability are well-documented (e.g., [Alschuler et al., 2013](#)). For people with SCI, prevalence rates of major depressive disorder are increased, and there are strong links between functional disability and depression ([Dryden et al., 2005](#)). Although successful treatment of depression and anxiety decrease disability levels (e.g., [Lenze et al., 2001](#)), the reverse association has also been well-established: Functional disability is related to lower levels of well-being (e.g., [Barry, Soulos, Murphy, Kasl, & Gill, 2013](#); [J. M. Hoffman, Bombardier, Graves, Kalpakjian, & Krause, 2011](#)). Longitudinal data have suggested that although people adjust to an injury in the year after it occurs and return close to previous levels of well-being (hedonic treadmill), they never quite return to pre-injury levels of life satisfaction (see [Lucas, 2007](#), for a review). Together, cross-sectional and longitudinal findings have shown that people who have sustained an SCI report lower levels of life

satisfaction and higher rates of depression compared with their noninjured counterparts, a pattern consistent with the results in the current study. What is unclear from previous literature, however, is whether age-related differences in affective experience documented in previous studies of relatively healthy people apply to people living with severe functional limitations.

Strength and Vulnerabilities of Aging

We predicted, based on SAVI, that age differences observed among people with no functional limitations would not replicate among people with SCI, particularly for affective reactivity—a dynamic measure that is believed to be indicative of emotion regulatory processes. SAVI predicts that strengths of aging lie in the ability of people to foresee negative events, and motivated by their time left in life (as posited by socioemotional selectivity; [Carstensen, 2006](#)) and their experiences gained throughout life, they proactively avoid these situations ([Charles & Luong, 2013](#)). When they cannot either avoid or extricate themselves from highly aversive circumstances, however, the vulnerabilities of physiological aging are hypothesized to make this experience more difficult to modulate. Although we did not study participants' physiological responses, we examined affective reactivity in response to daily stressors. As predicted, the age-related decreases in negative affective reactivity observed in the NSDE II sample were not replicated in the sample with SCI. In the sample with SCI, older adults were just as reactive to daily stressors as were younger adults.

Health, Age, and Emotion

A number of studies have examined the associations between age, health, and affect (e.g., [Ong, Mroczek, & Riffin, 2011](#)). Many of these studies have found that age is related to similar or lower levels of NA only after adjusting for physical health (e.g., [Kunzmann, Little, & Smith, 2000](#)). The current study varies from these prior investigations because it examined age differences in emotional experience among a group of people with functional disabilities. In prior studies, health problems were predominantly concentrated among older adults in the sample (e.g., [Piazza et al., 2007](#)). This skewed age distribution accurately reflects the greater prevalence of health problems in later life, but asks a somewhat different question than that in the current article. Our goal was to examine how patterns of age differences in daily emotional experiences and responses to daily stressors may vary for groups of people who differ on disability status.

Age, Emotional Well-Being, and Context

The current study builds on a growing number of studies that have examined age differences in well-being among samples of people who are faced with highly distressing situations, and do not find age-related benefits. For example, SAVI posits that a loss of social belonging represents a life context in which older adults lose age-related advantages in affective well-being (e.g., [Hawkey & Cacioppo, 2007](#)). In line with these predictions, one study showed that, in response to a social evaluative stressor, lonely older adults exhibited higher levels of reactivity and dampened recovery than their nonlonely and younger counterparts ([Ong, Rothstein, & Uchino, 2012](#)). Moreover, SAVI predicts that, in the face of

uncontrollable, chronic, or severe stressors, patterns of age differences in affective responses favoring older age may be attenuated or reversed. Consistent with this interpretation, an experience sampling study showed that when older individuals were confronted with complex hassles that influenced multiple life domains, they fared similar to, or even worse than, younger individuals in their affective responses to hassles (Wrzus, Müller, Wagner, Lindenberg, & Riediger, 2013). In contrast, older age was associated with less affective reactivity to hassles when dealing with circumscribed hassles that only affected one life domain, which presumably requires fewer coping efforts and resources.

Limitations and Future Directions

Life context offers one explanation for our findings showing that people with SCI do not display age differences in affective experience. Normative aging is associated with changes in life contexts that benefit daily affective well-being (e.g., Brose, Scheibe, & Schmiedek, 2013; Charles et al., 2010; Luong et al., 2011). For example, older adults are more likely to be retired, finished with child rearing, and, barring ill-health, have fewer constraints on how they spend their day than their younger counterparts who may have more competing demands on their time. Living with SCI, however, may nullify these age-related advantages, given that the condition is often accompanied by disruptions in daily routines, social networks, and resource capacities to cope with stressors (e.g., McColl & Rosenthal, 1994). The current study, however, did not study more microlevel processes to identify differences in these daily demands.

The study also did not include potential mediating mechanisms (e.g., physiological reactivity; use of specific emotion regulation strategies) that may account for the observed differences in age-related patterns between the SCI and NSDE samples. For example, SAVI would predict that one reason why older participants with SCI report lower affective well-being than their older, noninjured counterparts is that they are less able to effectively disengage from a caustic situation. Without directly asking people if they actively disengaged from a stressor, however, we are unable to adequately test this possibility. Thus, future research would benefit from a thorough examination of the types of emotion regulation strategies people living in different life contexts employ when faced with adverse situations.

It is also important to note that the use of cross-sectional data prevents us from ruling out the possibility that preexisting sample differences may account for the current findings. For example, it may be the case that even prior to their injury, the sample with SCI already exhibited lower levels of affective well-being than the non-SCI sample. Developmental trajectories also cannot be tested without longitudinal studies. Future studies should include prospective longitudinal designs to account for possible individual differences in affective experiences prior to injury, and to examine group differences in trajectories of affective experience and plausible mediating mechanisms therein.

Moreover, despite the matching process, the two samples varied on many demographic variables. In response, we adjusted for relevant group difference variables in our analyses, and matched for the most pertinent variables. One thing we were unable to assess, however, was possible cognitive impairment, because this was not assessed in either sample. Although individuals with brain

injuries were excluded from participating in the study, we cannot ignore the possibility that cognitive functioning may have played a role in our findings. Finally, whereas the non-SCI sample completed eight daily diary assessments, the sample with SCI completed 15 days of interviews in order to provide sufficient statistical power given the smaller sample size. Thus, selection biases could have contributed to these differences whereby younger and more highly educated individuals with SCI may have been more motivated to participate in the study. It is notable, however, that even if such selection biases were present in the current study, the sample with SCI still failed to show the same age-related advantages in affective experiences as the non-SCI sample.

Conclusion

The current study compared two national daily diary datasets of participants spanning a wide age range and who differed according to life context—namely, significant disability—which is posited to influence age-related patterns of affective experience in daily life. Our study makes important advances in the literature by testing theoretical predictions from SAVI in novel ways to understand under which circumstances we may (or may not) expect to observe age-related advantages in daily affective experiences and responses to stressors. We found that, in a situation where resources are taxed and older adults may be unable to leverage the emotion regulatory strengths of aging (i.e., in the context of severe functional disability), age-related advantages were attenuated or non-existent. The results bolster support for SAVI and provide a more comprehensive understanding of aging with a disability and its implications for affective well-being. Moreover, they reveal that perhaps the best way to assist older adults in difficult, chronically stressful situations is by providing environmental supports that would allow them to limit their exposure to aversive situations as opposed to focusing on cognitive strategies of emotion regulation.

References

- Aldwin, C. M., Jeong, Y.-J., Igarashi, H., Choun, S., & Spiro, A., III. (2014). Do hassles mediate between life events and mortality in older men?: Longitudinal findings from the VA Normative Aging Study. *Experimental Gerontology, 59*, 74–80. <http://dx.doi.org/10.1016/j.exger.2014.06.019>
- Aldwin, C. M., Sutton, K. J., Chiara, G., & Spiro, A., III. (1996). Age differences in stress, coping, and appraisal: Findings from the Normative Aging Study. *The Journals of Gerontology: Series B, Psychological Sciences and Social Sciences, 51B*, P179–P188. <http://dx.doi.org/10.1093/geronb/51B.4.P179>
- Almeida, D. M. (2005). Resilience and vulnerability to daily stressors assessed via diary methods. *Current Directions in Psychological Science, 14*, 64–68.
- Almeida, D. M., & Horn, M. C. (2004). Is daily life more stressful during middle adulthood? In O. G. Brim, C. D. Ryff, & R. C. Kessler (Eds.), *How healthy are we? A national study of well-being at midlife* (pp. 425–451). Chicago, IL: University of Chicago Press.
- Almeida, D. M., Wethington, E., & Kessler, R. C. (2002). The daily inventory of stressful events: An interview-based approach for measuring daily stressors. *Assessment, 9*, 41–55. <http://dx.doi.org/10.1177/1073191102009001006>
- Alschuler, K. N., Jensen, M. P., Sullivan-Singh, S. J., Borson, S., Smith, A. E., & Molton, I. R. (2013). The association of age, pain, and fatigue with physical functioning and depressive symptoms in persons with

- spinal cord injury. *The Journal of Spinal Cord Medicine*, 36, 483–491. <http://dx.doi.org/10.1179/2045772312Y.0000000072>
- Barry, L. C., Soulos, P. R., Murphy, T. E., Kasl, S. V., & Gill, T. M. (2013). Association between indicators of disability burden and subsequent depression among older persons. *The Journals of Gerontology: Series A, Biological Sciences and Medical Sciences*, 68, 286–292. <http://dx.doi.org/10.1093/gerona/gls179>
- Birditt, K. S., & Fingerma, K. L. (2005). Do we get better at picking our battles? Age group differences in descriptions of behavioral reactions to interpersonal tensions. *The Journals of Gerontology: Series B, Psychological Sciences and Social Sciences*, 60, P121–P128. <http://dx.doi.org/10.1093/geronb/60.3.P121>
- Birditt, K. S., Fingerma, K. L., & Almeida, D. M. (2005). Age differences in exposure and reactions to interpersonal tensions: A daily diary study. *Psychology and Aging*, 20, 330–340. <http://dx.doi.org/10.1037/0882-7974.20.2.330>
- Blanchard-Fields, F. (2007). Everyday problem solving and emotion: An adult developmental perspective. *Current Directions in Psychological Science*, 16, 26–31. <http://dx.doi.org/10.1111/j.1467-8721.2007.00469.x>
- Brose, A., Scheibe, S., & Schmiedek, F. (2013). Life contexts make a difference: Emotional stability in younger and older adults. *Psychology and Aging*, 28, 148–159. <http://dx.doi.org/10.1037/a0030047>
- Cacioppo, J. T., Berntson, G. G., Bechara, A., Tranel, D., & Hawkey, L. C. (2011). Could an aging brain contribute to subjective well-being? The value added by a social neuroscience perspective. In A. Todorov, S. Fiske, & D. Prentice (Eds.), *Social neuroscience: Toward understanding the underpinnings of the social mind* (pp. 249–262). New York, NY: Oxford University Press. <http://dx.doi.org/10.1093/acprof:oso/9780195316872.003.0017>
- Carstensen, L. L. (2006). The influence of a sense of time on human development. *Science*, 312, 1913–1915. <http://dx.doi.org/10.1126/science.1127488>
- Carstensen, L. L., Turan, B., Scheibe, S., Ram, N., Ersner-Hershfield, H., Samanez-Larkin, G. R., . . . Nesselroade, J. R. (2011). Emotional experience improves with age: Evidence based on over 10 years of experience sampling. *Psychology and Aging*, 26, 21–33. <http://dx.doi.org/10.1037/a0021285>
- Charles, S. T. (2010). Strength and vulnerability integration: A model of emotional well-being across adulthood. *Psychological Bulletin*, 136, 1068–1091. <http://dx.doi.org/10.1037/a0021232>
- Charles, S. T., & Carstensen, L. L. (2010). Social and emotional aging. *Annual Review of Psychology*, 61, 383–409. <http://dx.doi.org/10.1146/annurev.psych.093008.100448>
- Charles, S. T., & Luong, G. (2013). Emotional experience across adulthood: The theoretical model of strength and vulnerability integration. *Current Directions in Psychological Science*, 22, 443–448. <http://dx.doi.org/10.1177/0963721413497013>
- Charles, S. T., Luong, G., Almeida, D. M., Ryff, C., Sturm, M., & Love, G. (2010). Fewer ups and downs: Daily stressors mediate age differences in negative affect. *The Journals of Gerontology: Series B, Psychological Sciences and Social Sciences*, 65B, 279–286. <http://dx.doi.org/10.1093/geronb/gbq002>
- Charles, S. T., & Piazza, J. R. (2009). Age differences in affective well-being: Context matters. *Social and Personality Psychology Compass*, 3, 711–724. <http://dx.doi.org/10.1111/j.1751-9004.2009.00202.x>
- Charles, S. T., Piazza, J. R., Luong, G., & Almeida, D. M. (2009). Now you see it, now you don't: Age differences in affective reactivity to social tensions. *Psychology and Aging*, 24, 645–653. <http://dx.doi.org/10.1037/a0016673>
- Degraff, A. H., & Schaffer, J. (2008). Emotion-focused coping: A primary defense against stress for people living with spinal cord injury. *Journal of Rehabilitation*, 74, 19–24.
- Diehl, M., & Hay, E. L. (2010). Risk and resilience factors in coping with daily stress in adulthood: The role of age, self-concept incoherence, and personal control. *Developmental Psychology*, 46, 1132–1146. <http://dx.doi.org/10.1037/a0019937>
- Dryden, D. M., Saunders, L. D., Rowe, B. H., May, L. A., Yiannakoulis, N., Svenson, L. W., . . . Voaklander, D. C. (2005). Depression following traumatic spinal cord injury. *Neuroepidemiology*, 25, 55–61. <http://dx.doi.org/10.1159/000086284>
- Fuhrer, M. J. (1996). The subjective well-being of people with spinal cord injury: Relationships to impairment, disability and handicap. *Topics in Spinal Cord Injury Rehabilitation*, 1, 56–71.
- Garshick, E., Kelley, A., Cohen, S. A., Garrison, A., Tun, C. G., Gagnon, D., & Brown, R. (2005). A prospective assessment of mortality in chronic spinal cord injury. *Spinal Cord*, 43, 408–416. <http://dx.doi.org/10.1038/sj.sc.3101729>
- Gerstorff, D., Ram, N., Mayraz, G., Hidajat, M., Lindenberg, U., Wagner, G. G., & Schupp, J. (2010). Late-life decline in well-being across adulthood in Germany, the United Kingdom, and the United States: Something is seriously wrong at the end of life. *Psychology and Aging*, 25, 477–485. <http://dx.doi.org/10.1037/a0017543>
- Giannini, M. J., Bergmark, B., Kreshover, S., Elias, E., Plummer, C., & O'Keefe, E. (2010). Understanding suicide and disability through three major disabling conditions: Intellectual disability, spinal cord injury, and multiple sclerosis. *Disability and Health Journal*, 3, 74–78. <http://dx.doi.org/10.1016/j.dhjo.2009.09.001>
- Hawkey, L. C., & Cacioppo, J. T. (2007). Aging and loneliness: Downhill quickly? *Current Directions in Psychological Science*, 16, 187–191. <http://dx.doi.org/10.1111/j.1467-8721.2007.00501.x>
- Hoffman, J. M., Bombardier, C. H., Graves, D. E., Kalpakjian, C. Z., & Krause, J. S. (2011). A longitudinal study of depression from 1 to 5 years after spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 92, 411–418. <http://dx.doi.org/10.1016/j.apmr.2010.10.036>
- Hoffman, L., & Stawski, R. S. (2009). Persons as context: Evaluating between-person and within-person effects in longitudinal analysis. *Research in Human Development*, 6, 97–120. <http://dx.doi.org/10.1080/15427600902911189>
- Isaacowitz, D. M., & Blanchard-Fields, F. (2012). Linking process and outcome in the study of emotion and aging. *Perspectives on Psychological Science*, 7, 3–17. <http://dx.doi.org/10.1177/1745691611424750>
- Kemp, B. J., & Krause, J. S. (1999). Depression and life satisfaction among people ageing with post-polio and spinal cord injury. *Disability and Rehabilitation*, 21, 241–249. <http://dx.doi.org/10.1080/096382899297666>
- Kessler, R. C., Andrews, G., Colpe, L. J., Hiripi, E., Mroczek, D. K., Normand, S.-L. T., . . . Zaslavsky, A. M. (2002). Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological Medicine*, 32, 959–976. <http://dx.doi.org/10.1017/S0033291702006074>
- Krause, J. S. (1998). Changes in adjustment after spinal cord injury: A 20-year longitudinal study. *Rehabilitation Psychology*, 43, 41–55. <http://dx.doi.org/10.1037/0090-5550.43.1.41>
- Krause, J. S., & Crewe, N. M. (1991). Chronologic age, time since injury, and time of measurement: Effect on adjustment after spinal cord injury. *Archives of Physical Medicine and Rehabilitation*, 72, 91–100.
- Kunzmann, U., Little, T. D., & Smith, J. (2000). Is age-related stability of subjective well-being a paradox? Cross-sectional and longitudinal evidence from the Berlin Aging Study. *Psychology and Aging*, 15, 511–526. <http://dx.doi.org/10.1037/0882-7974.15.3.511>
- Labouvie-Vief, G. (2003). Dynamic integration: Affect, cognition, and the self in adulthood. *Current Directions in Psychological Science*, 12, 201–206. <http://dx.doi.org/10.1046/j.0963-7214.2003.01262.x>
- Leary, M. (1990). Responses to social exclusion: Social anxiety, jealousy, loneliness, depression, and low self-esteem. *Journal of Social and Clinical Psychology*, 9, 221–229. <http://dx.doi.org/10.1521/jscp.1990.9.2.221>

- Lenze, E. J., Rogers, J. C., Martire, L. M., Mulsant, B. H., Rollman, B. L., Dew, M. A., . . . Reynolds, C. F., III. (2001). The association of late-life depression and anxiety with physical disability: A review of the literature and prospectus for future research. *The American Journal of Geriatric Psychiatry*, 9, 113–135. <http://dx.doi.org/10.1097/00019442-200105000-00004>
- Lucas, R. E. (2007). Adaptation and the set-point model of subjective well-being: Does happiness change after major life events? *Current Directions in Psychological Science*, 16, 75–79. <http://dx.doi.org/10.1111/j.1467-8721.2007.00479.x>
- Luong, G., & Charles, S. T. (2014). Age differences in affective and cardiovascular responses to a negative social interaction: The role of goals, appraisals, and emotion regulation. *Developmental Psychology*, 50, 1919–1930. <http://dx.doi.org/10.1037/a0036621>
- Luong, G., Charles, S. T., & Fingerman, K. L. (2011). Better with age: Social relationships across adulthood. *Journal of Social and Personal Relationships*, 28, 9–23. <http://dx.doi.org/10.1177/0265407510391362>
- Lysack, C., Komanecy, M., Kabel, A., Cross, K., & Neufeld, S. (2007). Environmental factors and their role in community integration after spinal cord injury. *Canadian Journal of Occupational Therapy/Revue Canadienne d'Ergothérapie*, 74, 243–254. <http://dx.doi.org/10.1177/000841740707405503>
- Marmot, M., Ryff, C. D., Bumpass, L. L., Shipley, M., & Marks, N. F. (1997). Social inequalities in health: Next questions and converging evidence. *Social Science & Medicine*, 44, 901–910. [http://dx.doi.org/10.1016/S0277-9536\(96\)00194-3](http://dx.doi.org/10.1016/S0277-9536(96)00194-3)
- McCull, M. A., & Rosenthal, C. (1994). A model of resource needs of aging spinal cord injured men. *Paraplegia*, 32, 261–270. <http://dx.doi.org/10.1038/sc.1994.46>
- Migliorini, C., Tonge, B., & Taleporos, G. (2008). Spinal cord injury and mental health. *Australian and New Zealand Journal of Psychiatry*, 42, 309–314. <http://dx.doi.org/10.1080/00048670801886080>
- Mroczek, D. K. (2001). Age and emotion in adulthood. *Current Directions in Psychological Science*, 10, 87–90. <http://dx.doi.org/10.1111/1467-8721.00122>
- National Spinal Cord Injury Statistical Center. (2013 February). *Spinal cord injury facts and figures at a glance*. Birmingham, AL: Author. Retrieved from https://www.nscisc.uab.edu/PublicDocuments/facts_figures_docs/Facts%202013.pdf
- Neugarten, B. L. (1979). Time, age, and the life cycle. *The American Journal of Psychiatry*, 136, 887–894. <http://dx.doi.org/10.1176/ajp.136.7.887>
- Neupert, S. D., Almeida, D. M., & Charles, S. T. (2007). Age differences in reactivity to daily stressors: The role of personal control. *The Journals of Gerontology: Series B, Psychological Sciences and Social Sciences*, 62, P216–P225. <http://dx.doi.org/10.1093/geronb/62.4.P216>
- Noreau, L., & Fougereyrollas, P. (2000). Long-term consequences of spinal cord injury on social participation: The occurrence of handicap situations. *Disability and Rehabilitation*, 22, 170–180. <http://dx.doi.org/10.1080/096382800296863>
- Ong, A. D., Mroczek, D. K., & Riffin, C. (2011). The health significance of positive emotions in adulthood and later life. *Social and Personality Psychology Compass*, 5, 538–551. <http://dx.doi.org/10.1111/j.1751-9004.2011.00370.x>
- Ong, A. D., Rothstein, J. D., & Uchino, B. N. (2012). Loneliness accentuates age differences in cardiovascular responses to social evaluative threat. *Psychology and Aging*, 27, 190–198. <http://dx.doi.org/10.1037/a0025570>
- Park, D. C. (2002). Aging, cognition, and culture: A neuroscientific perspective. *Neuroscience and Biobehavioral Reviews*, 26, 859–867. [http://dx.doi.org/10.1016/S0149-7634\(02\)00072-6](http://dx.doi.org/10.1016/S0149-7634(02)00072-6)
- Piazza, J. R., Charles, S. T., & Almeida, D. M. (2007). Living with chronic health conditions: Age differences in affective well-being. *Journals of Gerontology: Psychological Sciences*, 62, 313–321.
- Raudenbush, S. W., & Bryk, A. S. (2002). *Hierarchical linear models: Applications and data analysis methods*. Thousand Oaks, CA: Sage.
- Riediger, M., Schmiedek, F., Wagner, G. G., & Lindenberger, U. (2009). Seeking pleasure and seeking pain: Differences in prohedonic and contra-hedonic motivation from adolescence to old age. *Psychological Science*, 20, 1529–1535. <http://dx.doi.org/10.1111/j.1467-9280.2009.02473.x>
- Salthouse, T. A. (2004). What and when of cognitive aging. *Current Directions in Psychological Science*, 13, 140–144. <http://dx.doi.org/10.1111/j.0963-7214.2004.00293.x>
- Schopp, L. H., Clark, M. J., Hagglund, K. J., Sherman, A. K., Stout, B. J., Gray, D. B., & Boninger, M. L. (2007). Life activities among individuals with spinal cord injury living in the community: Perceived choice and perceived barriers. *Rehabilitation Psychology*, 52, 82–88. <http://dx.doi.org/10.1037/0090-5550.52.1.82>
- Sliwinski, M. J., Almeida, D. M., Smyth, J., & Stawski, R. S. (2009). Intraindividual change and variability in daily stress processes: Findings from two measurement-burst diary studies. *Psychology and Aging*, 24, 828–840. <http://dx.doi.org/10.1037/a0017925>
- Stone, A. A., Schwartz, J. E., Broderick, J. E., & Deaton, A. (2010). A snapshot of the age distribution of psychological well-being in the United States. *PNAS: Proceedings of the National Academy of Sciences of the United States of America*, 107, 9985–9990. <http://dx.doi.org/10.1073/pnas.1003744107>
- Stroebe, M., Schut, H., & Stroebe, W. (2007). Health outcomes of bereavement. *The Lancet*, 370, 1960–1973. [http://dx.doi.org/10.1016/S0140-6736\(07\)61816-9](http://dx.doi.org/10.1016/S0140-6736(07)61816-9)
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54, 1063–1070. <http://dx.doi.org/10.1037/0022-3514.54.6.1063>
- Wolff, J. L., Starfield, B., & Anderson, G. (2002). Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *JAMA Internal Medicine*, 162, 2269–2276. <http://dx.doi.org/10.1001/archinte.162.20.2269>
- Wrzus, C., Luong, G., Wagner, G. G., & Riediger, M. (2015). Can't get it out of my head: Age differences in affective responsiveness vary with preoccupation and elapsed time after daily hassles. *Emotion*, 15, 257–269. <http://dx.doi.org/10.1037/emo0000019>
- Wrzus, C., Müller, V., Wagner, G. G., Lindenberger, U., & Riediger, M. (2013). Affective and cardiovascular responding to unpleasant events from adolescence to old age: Complexity of events matters. *Developmental Psychology*, 49, 384–397. <http://dx.doi.org/10.1037/a0028325>
- Yeo, J. D., Walsh, J., Rutkowski, S., Soden, R., Craven, M., & Middleton, J. (1998). Mortality following spinal cord injury. *Spinal Cord*, 36, 329–336. <http://dx.doi.org/10.1038/sj.sc.3100628>

Received September 29, 2014

Revision received June 30, 2015

Accepted July 5, 2015 ■