

The Oxford Handbook of Integrative Health Science

Carol D. Ryff (ed.), Robert F. Krueger (ed.)

<https://doi.org/10.1093/oxfordhb/9780190676384.001.0001>

Published online: 09 October 2018 **Published in print:** 29 November 2018

Online ISBN:

9780190676407

Print ISBN: 9780190676384

Search in this book

CHAPTER

4 Early Life Adversity and Adult Health

p. 49

Cynthia S. Levine, Gregory E. Miller, Margie E. Lachman, Teresa E. Seeman, Edith Chen

<https://doi.org/10.1093/oxfordhb/9780190676384.013.3> Pages 49–62

Published: 09 October 2018

Abstract

Research has shown that early life adversity can have implications for health later in life. Specifically, socioeconomic disadvantage, parental maltreatment, and parent divorce and death in childhood have been linked to cardiovascular disease, diabetes, cancer, and mortality in adulthood. Increasingly, recent research has focused on which factors can protect against these poor health outcomes and what promotes resilience, despite early life adversity. This chapter reviews research linking early life adversity to health, with a focus on highlighting the psychosocial factors that play this type of protective role. These factors include social and relational ones, such as maternal nurturance, as well as beliefs and coping strategies. The chapter concludes by suggesting areas of future research, including additional investigation of which psychosocial factors protect health, how multiple psychosocial factors might interact to protect health, and how early life adversity might affect adult health across different groups throughout the life span.

Keywords: [early life adversity](#), [resilience](#), [psychosocial factors](#), [social factors](#), [relational factors](#), [maternal nurturance](#), [beliefs](#), [coping strategies](#), [adult health](#), [health outcomes](#)

Subject: [Health Psychology](#), [Psychology](#)

Series: [Oxford Library of Psychology](#)

Collection: [Oxford Handbooks Online](#)

Introduction

A large body of literature suggests that psychosocial stress in early childhood has implications for a range of health outcomes later in life (Miller, Chen, & Parker, 2011; Repetti, Taylor, & Seeman, 2002; Shonkoff, Boyce, & McEwen, 2009; Wegman & Stetler, 2009). Growing up in lower socioeconomic status (SES) environments, being mistreated in childhood, or experiencing other types of childhood adversity increases one's risk for a range of chronic diseases of aging and earlier mortality. Previous literature has demonstrated that these associations can be explained in part by a number of factors, including biological programming of pro-inflammatory tendencies and hormone dysregulation that are initiated during childhood, unhealthy behaviors, and reduced physical and psychosocial resources, among others (Braveman & Barclay, 2009; Danese & McEwen, 2012; Repetti, Robles, & Reynolds, 2011; Shonkoff & Garner, 2012; Taylor, 2010). Recently, however, research has focused increasingly on which factors can protect those who experience early life adversity from poor health outcomes (Chen & Miller, 2012).

The present chapter reviews the literature on the association between early life adversity and adult health outcomes, focusing in particular on the psychosocial factors that may protect the health of people who have experienced adversity in early life. We begin by reviewing the literature on early life adversity and adult health, which shows that people who have grown up in lower SES environments, people who have experienced abuse as children, and others who have experienced adversity have a range of poor health outcomes, including greater risk for cardiovascular disease, cancer, Type 2 diabetes, and mortality. We then briefly review some mechanisms underlying this association. Next, we review research showing which psychosocial factors might buffer people who have experienced adversity against poor health outcomes. These include social relationships, especially a warm family environment and maternal nurturance, as well as certain strategies for coping with adversity. We conclude by outlining directions for future research. Throughout the chapter, we focus primarily, although not exclusively, on studies from the Midlife in the United States (MIDUS) study. Because MIDUS assesses a wide range of psychosocial factors, including multiple types of early life adversity, and includes biomarker data, it is well positioned to illuminate multiple factors that contribute to or mitigate the associations between early life adversity and later life health.

p. 50

Background Literature on Early Life Adversity and Adult Health

A robust association between early life adversity and adult health has been found across a range of types of adversity and health outcomes. With respect to self-reported outcomes, research using samples of adults in MIDUS, other samples of healthy middle-aged adults, and healthy children showed that people who have experienced different types of adversities in childhood, including poverty and other economic difficulties, exposure to violence, and frequent moves, all rated their overall health as worse in adulthood (Boynton-Jarrett, Ryan, Berkman, & Wright, 2008; Bures, 2003; Greenfield & Marks, 2009a; Laaksonen, Rahkonen, Martikainen, & Lahelma, 2005). Research using MIDUS further showed that adversity in early life also predicts later self-reported functional limitations, such as difficulty climbing up a flight of stairs, and a higher number of acute symptoms and chronic conditions in adulthood (Greenfield & Marks, 2009a; Maier & Lachman, 2000).

Similar early life experiences have also been linked to physiological measures, such as metabolic syndrome, allostatic load, and inflammation, which indicate heightened risk for a number of chronic diseases of aging. For example, research from MIDUS showed that people who grow up in lower SES environments are more likely to develop metabolic syndrome in adulthood (Miller, Lachman, et al., 2011). This is a cluster of symptoms (defined by the International Diabetes Federation as including abdominal obesity, as well as

elevated blood pressure, elevated triglycerides, elevated fasting glucose, and low high-density lipoprotein levels; Cornier et al., 2008) that is associated with risk for cardiovascular disease, Type 2 diabetes, and all-cause mortality (Cornier et al., 2008; Ford, 2005; Lakka et al., 2002). A similar association has been documented in MIDUS between self-reported abuse in childhood and metabolic syndrome in adulthood (Lee, Tsenkova, & Carr, 2014). MIDUS data also show that people who have experienced SES disadvantage or other stressors (e.g., parental death, divorce, or abuse) in early life are at greater risk for developing higher levels of allostatic load, or the dysregulation that develops across multiple systems as the result of repeated attempts to maintain allostasis in the face of ongoing stress (Friedman, Karlamangla, Gruenewald, Koretz, & Seeman, 2015; Gruenewald et al., 2012). Allostatic load, in turn, has been linked to conditions such as hypertension, obesity, diabetes, and cardiovascular disease (Juster, McEwen, & Lupien, 2010; Seeman, Epel, Gruenewald, Karlamangla, & McEwen, 2010; Seeman, McEwen, Rowe, & Singer, 2001). Finally, early life adversity has been linked to higher levels of inflammation in adults in MIDUS (Hostinar, Lachman, Mroczek, Seeman, & Miller, 2015; Slopen et al., 2010), with inflammatory markers in turn predicting risk for cardiovascular disease and other chronic diseases of aging (Ridker, Hennekens, Buring, & Rifai, 2000; Ridker, Rifai, Stampfer, & Hennekens, 2000).

As these associations between early life adversity and physiological risk suggest, early life adversity is further associated with risk for the development of chronic diseases of aging and, ultimately, mortality. Studies using MIDUS (Friedman, Montez, Sheehan, Gruenewald, & Seeman, 2015) and other data (Dong et al., 2004) showed that experiencing a greater number of different types of adverse events in childhood predicted cardiovascular disease in adulthood. Early life adversity has similarly been linked to cancer. For instance, a study of Jewish Israelis who emigrated from Europe after World War II (i.e., had potentially been exposed to the Holocaust) were found to have higher rates of cancer than those who left earlier (Keinan-Boker, Vin-Raviv, Liphshitz, Linn, & Barchana, 2009). Abuse in childhood also predicts cancer occurrence in adult MIDUS participants (Morton, Schafer, & Ferraro, 2012). Finally, as this heightened risk for diseases such as cardiovascular disease and cancer might suggest, different types of early life adversity have also been linked to mortality. Research using MIDUS data has found that women who report being abused in childhood have higher rates of all-cause mortality, even controlling for depression (Chen, Turiano, Mroczek, & Miller, 2016). Research using other data has also linked higher mortality rates to lower childhood SES (Galobardes, Lynch, & Davey Smith, 2004) and to experiencing a higher number of adverse events in childhood (Kelly-Irving et al., 2013).

p. 51

Mechanisms: MIDUS Advances on Health Behaviors and Psychological Resources

Research, including work using MIDUS, has shed light on some of the psychosocial and behavioral mechanisms through which early life adversity might give rise to poor health in adulthood. While there are a number of biological and psychosocial mechanisms thought to underlie the association between early life adversity and adult health, we focus here on psychosocial factors that have been highlighted by research with MIDUS, namely health behaviors and psychological resources.

Early life adversity is associated with a number of unhealthy behaviors in adulthood, as shown by multiple studies using MIDUS data. For example, those who experience physical abuse in childhood are more likely to smoke regularly as adults and to smoke for a longer period of time, which partially explains their greater incidence of respiratory disease (Goodwin & Wamboldt, 2012; Taha, Galea, Hien, & Goodwin, 2014). Having experienced psychological and physical violence in childhood is also linked to a greater likelihood of using food to cope with stress (Greenfield & Marks, 2009b). In turn, this way of coping with stress has been linked to the consumption of high-fat and high-sugar foods, obesity, and poor health in samples of healthy adults (i.e., medical students, adults from the community who came into the lab; Epel et al., 2004; Oliver, Wardle,

& Gibson, 2000; Tsenkova, Boylan, & Ryff, 2012). Use of food to cope with stress also explains part of the association between childhood violence and obesity in adulthood (Greenfield & Marks, 2009b). Finally, childhood socioeconomic disadvantage and other adverse childhood experiences predict decreased physical activity in adulthood, which is one of the mediators of the relationship between early childhood adversity and adverse outcomes, such as higher levels of inflammation and diabetes in adulthood (Hostinar et al., 2015; Tsenkova, Pudrovska, & Karlamangla, 2014).

In addition to unhealthy behaviors, research using MIDUS shows that early life adversity has been linked to psychological states that give rise to poor health in adulthood. The majority of this work with MIDUS has focused on the role of psychological well-being. For example, research using MIDUS data showed that early life adversity can increase one's risk for depressive symptoms, negative affective states, or other indices of lower psychological well-being in adulthood. Specifically, people experiencing adversities in childhood, such as parental divorce or having parents who used authoritarian (high demands and low responsiveness) rather than authoritative (more reasonable demands and high responsiveness) styles of parenting, report more depressive symptoms as adults (Maier & Lachman, 2000; Rothrauff, Cooney, & An, 2009; Uphold-Carrier & Utz, 2012). Furthermore, people who experienced abuse as children have higher levels of distress, including depression and anxiety, as adults (Schrepf, Markon, & Lutgendorf, 2014). Childhood psychological violence and lower quality parental relationships predict lower frequency of positive affect and higher frequency of negative affect in adulthood (Greenfield & Marks, 2010b; Mellers, Charles, Neupert, & Almeida, 2010). Finally, people who experienced a lack of affection and support from parents or abuse from parents in childhood have lower levels of eudaimonic psychological well-being (i.e., a sense a meaning or self-realization) as adults (An & Cooney, 2006; Greenfield & Marks, 2010a). All of these negative psychological states are, in turn, linked to worse physiological and health outcomes, such as higher levels of inflammation and increased risk of cardiovascular disease in MIDUS and other samples of healthy adults (Friedman, Hayney, Love, Singer, & Ryff, 2007; Kubzansky & Kawachi, 2000; Pressman & Cohen, 2005; Ryff, Singer, & Love, 2004; Schrepf et al., 2014).

Moderators: Social Relationships

Although early life adversity increases one's risk for poor health in adulthood, not everyone who experiences adversity in childhood gets sick. Recent research has focused on identifying protective factors that help to buffer against the health risk of early life adversity. We review two types of buffers that research using MIDUS, as well as other samples, has identified: social relationships and individual psychological traits that help people to cope with adversity (Figure 4.1).

A large body of literature highlights the role that positive social relationships and support can play in fostering good health outcomes (see Cohen, 2004; Uchino, Cacioppo, & Kiecolt-Glaser, 1996, for reviews). For children who experience adversity, one type of social relationship that has the potential to play an especially important role in protecting health down the line is a positive and supportive family environment. In particular, research has focused on nurturant parenting, especially from mothers. Nurturing parents who are warm and sensitive have been shown to provide a variety of educational and psychological advantages to children who experience adversity (see Luthar, 2006; Masten, 2001, for reviews). For example, having nurturant parents helps children to feel that the world is a safe place and helps them to develop adaptive emotion regulation strategies (see Cassidy & Shaver, 2008; Repetti et al., 2002, for reviews). Building on the work highlighting such psychological benefits, a number of studies have investigated the role that nurturant parenting can play in protecting the health of those who experience adversity in early life.




Figure 4.1 Effect of early life adversity on adult health is moderated by social relationships and by individual traits, such as adaptive coping strategies.

With its national sample and depth and breadth of psychosocial variables, in addition to biological and health data, MIDUS is especially well positioned to help researchers identify which social relationship factors buffer against the deleterious health implications of early life adversity. Accordingly, researchers have used MIDUS data to explore family relationships—especially relationships with parents—as one pathway to resilience in the wake of early life adversities such as lower early life SES and physical and emotional abuse in childhood. With regard to early life SES, one study using MIDUS data investigated whether parental nurturance might play a role in protecting the health of those who grew up in lower SES environments (Miller, Lachman, et al., 2011). These authors used participants from the MIDUS biomarker sample, a subset of over a thousand MIDUS participants who traveled to a General Clinical Research Center (GCRC) for an overnight visit and from whom biomarker data were collected.

Consistent with previous research, participants' childhood SES (here, their parents' level of educational attainment) was related to participants' metabolic syndrome symptoms. As noted, this is a cluster of symptoms (abdominal obesity, as well as elevated blood pressure, elevated triglycerides, elevated fasting glucose, and low high-density lipoprotein levels; Cornier et al., 2008) that is associated with risk for developing cardiovascular disease, Type 2 diabetes, and all-cause mortality (Cornier et al., 2008; Ford, 2005; Lakka et al., 2002). However, this association was moderated by participants' relationships with their parents. Although people whose parents had lower levels of educational attainment had a higher number of metabolic syndrome symptoms overall, this effect was less pronounced among those who reported that their mothers had been more nurturing (e.g., understood their problems and worries, gave them attention when they needed it). Indeed, among those with the most nurturing mothers, there was no relationship between early life SES and metabolic syndrome. Such a buffering effect did not emerge for paternal nurturance, which the authors suggested may result from the adults in MIDUS largely being of a generation when mothers, rather than fathers, were largely responsible for childrearing.

p. 53

These findings are consistent with other non-MIDUS data showing that positive relationships with family members can protect children from the negative physiological correlates of early life SES. For example, in samples of middle school children and healthy adults from the community, maternal warmth has been found to protect children who grew up in lower SES environments against higher levels of allostatic load and pro-inflammatory profiles that would typically be characteristic of this group (Chen, Miller, Kobor, & Cole, 2011; Evans, Kim, Ting, Teshler, & Shannis, 2007). Experimental evidence lends support to the idea that better relationships with parents play a causal role in protecting children biologically when they experience

adversity. Among a sample of African Americans in the rural southern United States who were largely from lower SES families, those who were randomly assigned to a parenting intervention that taught nurturant-involved parenting skills had children who displayed lower levels of low-grade inflammation 8 years later compared to those who were in a control group (Miller, Brody, Yu, & Chen, 2014).

Positive parental relationships during childhood can buffer against the negative health consequences of emotional and physical abuse in childhood as well. In a study using MIDUS participants, Schafer, Morton, and Ferraro (2014) found that, although people who had been abused in childhood had worse self-reported health, more frequent physical symptoms, and a higher number of chronic conditions in adulthood, this relationship was attenuated when they also rated their relationship with both parents as being “good,” “very good,” or “excellent” overall when they were growing up. The authors suggested that these findings may emerge because the children who had positive relationships with their parents are able to use those relationships as a foundation to forgive or reconcile with their parents as they grow older.

These results are consistent with other non-MIDUS research showing that, although having a harsh relationship with one parent is typically associated with declines in self-rated health over time among adolescents, this relationship is attenuated among adolescents (i.e., seventh graders who were part of a larger study on economic hardship and health) who have a warm relationship with the other parent (Schofield, Conger, Gonzales, & Merrick, 2016). A similar pattern emerges with physiological outcomes. Research using adult participants in the Coronary Artery Risk Development in Young Adults (CARDIA) study further showed that abuse in childhood predicted levels of allostatic load among people whose parents were less warm and affectionate, but not among those whose parents were more warm and affectionate (Carroll et al., 2013). Again, experimental evidence suggests that improving parenting can play a causal role in ameliorating the physiological outcomes of children who have experienced mistreatment. For example, teaching parenting skills to foster parents results in greater declines in daily cortisol levels among children who have been mistreated (Fisher, Gunnar, Chamberlain, & Reid, 2000). Thus, across multiple types of early life adversity, positive relationships with one’s parents consistently emerge as a protective factor for physiological and health outcomes across the life course.

Although the MIDUS research on relationships that can buffer children against childhood adversity has focused on parental relationships, some additional studies use other data to show that a wider range of types of social connection, including those with peers, adults outside the home, or the community more broadly, can have a similarly protective effect. For example, one study of African American youth in the rural southern United States has shown that emotional support—defined broadly as a composite of emotional support from parents, peers, and other adult mentors—can buffer against the physiological correlates of childhood socioeconomic adversity (Brody, Lei, Chen, & Miller, 2014). Specifically, these authors examined allostatic load levels at age 19 as a function of changes in neighborhood poverty across adolescence. Allostatic load was higher among individuals who lived in progressively poorer neighborhoods throughout their adolescence, relative to those who lived in neighborhoods with consistent levels of poverty or who shifted from neighborhoods with higher poverty to neighborhoods with lower poverty. However, even among those who lived in progressively poorer neighborhoods, emotional support played a buffering role. Increasing neighborhood poverty from childhood to adolescence was associated with greater allostatic load among those who had lower levels of emotional support from parents, peers, and other mentors, but not among those who had higher levels of emotional support.

p. 54 In addition, another study found that supportive role models, or adult figures the children admires and wishes to emulate or sees as providing support, can protect lower SES children against poor physiological outcomes (Chen, Lee, Cavey, & Ho, 2013). Specifically, among children who do not have supportive role models, lower SES (family savings) predicted higher levels of the pro-inflammatory cytokine interleukin 6 (IL-6). However, when lower SES children identified a supportive role model, their IL-6 levels were comparable to those of their higher SES counterparts.

Similarly, neighborhood social relationships can also protect the health of children who experience adversity. Specifically, the social capital of the community in which a child lives (e.g., the extent to which parents and their children can trust others in the community and are integrated into the community) can buffer against the effects of childhood adversity. Another study, again using data other than MIDUS, found that children in families with lower incomes were more likely to smoke and had a higher body mass index (BMI) when they lived in communities with lower levels of social capital, which was defined here as a less cohesive community, a community where adults were less likely to intervene to protect children, and a community where children had less supportive relationships with adults. However, among children living in communities with high social capital, having lower family income did not increase the likelihood of the child smoking or having a high BMI (Evans & Kutcher, 2011). Similarly, research suggested that children whose families lived in poverty were more likely to smoke if they lived in communities where people did not know each other, where parents were not involved in school and community organizations, and where children were not involved in athletics. However, the link between poverty and smoking was weaker among children who lived in communities that did have such resources (Thorlindsson, Valdimarsdottir, & Hrafn, 2012). Finally, research using a non-MIDUS community sample showed that social trust, or the belief that others in the community are reliable, reduced metabolic risk among adults who grew up in lower SES circumstances (Hostinar, Ross, Chen, & Miller, 2017).

Moderators: Individual Psychological Traits

The majority of research on factors that protect against the negative health consequences of early life adversity has focused on relational factors. This includes research on this topic using MIDUS data. However, a smaller literature has begun to highlight the role that individual psychological traits, such as beliefs or strategies for coping with adversity, might also play in protecting the health of those who have experienced adversity in childhood. One example of such an individual trait that has received particular attention is a coping strategy known as “shift and persist” (Chen & Miller, 2012). This set of beliefs, which involves both (1) shifting or accepting stressors and as a result adjusting the self to the environment and (2) persisting or enduring and finding meaning or optimism despite adversity, has been found to protect the health of lower SES children. We next review research from MIDUS and other samples that highlights the protective role of shift-and-persist beliefs, along with research on other individual traits that might also protect the health of those who experience early life adversity.

Shift-and-Persist Coping Strategies

Across multiple studies, Chen and colleagues (Chen & Miller, 2012; Chen, Strunk, et al., 2011) have identified shift-and-persist coping strategies as protective for the health of lower SES children. The theory underlying these findings relies on the understanding that lower SES environments tend to have particular characteristics. Thus, the traits that allow children to cope effectively with the stressors that they are likely to encounter in such environments will be most effective at promoting resilience. Lower SES environments tend to be characterized by frequent, recurring, uncontrollable stressors (Brady & Matthews, 2002). Lower SES families also often face many competing demands and have limited material resources to deal with them. They are also likely to live in neighborhoods with higher rates of violence (Buka, Stichick, Birdthistle, & Earls, 2001). In such environments, where it is often not possible to influence or change the situation, it can be beneficial to shift or adjust oneself to the stressors. This can entail cognitive reappraisal of stressful situations or the use of other self-regulation strategies that involve changing one's interpretation of or response to the situation. At the same time, shifting does not entail giving up or losing hope for the future. Consistent with the trauma resilience literature showing that meaning and optimism can help people adapt to difficult experiences (Bonanno, 2004; Dunkel Schetter & Dolbier, 2011), the shift-and-persist model suggests that, in addition to shifting, it is adaptive to persist, or maintain a focus on long-term goals by finding meaning in life and maintaining optimism about the future.

p. 55

Research using samples of children from the community and adults from MIDUS supported the idea that shift-and-persist strategies are physiologically protective for people from lower SES backgrounds. For example, shift-and-persist strategies have been shown to protect healthy lower SES children against a higher BMI and higher levels of inflammation and lower SES children with asthma against heightened asthma-relevant inflammatory markers (Chen, McLean, & Miller, 2015; Chen, Strunk, et al., 2011; Kalleem et al., 2013). The same protective effect does not emerge for higher SES children. In addition, research using MIDUS participants found that, among those who grew up in lower SES circumstances, there was a significant two-way interaction of shift by persist in predicting allostatic load. In contrast, no such interaction emerged among those who grew up high in SES (Chen, Miller, Lachman, Gruenewald, & Seeman, 2012).

Other Individual Characteristics

There is relatively little research on other individual-level characteristics besides shift-and-persist strategies that might protect people who have experienced *early* life adversity in particular. However, a number of studies using data from MIDUS showed that other individual psychological characteristics might protect the health of those who are lower SES *in adulthood*. For example, among lower SES adults in MIDUS, a higher sense of control is linked to better self-rated health, fewer acute symptoms and functional limitations, and lower mortality rates (Lachman & Weaver, 1998; Turiano, Chapman, Agrigoroaei, Infurna, & Lachman, 2014). In addition, higher psychological well-being buffers individuals with lower levels of educational attainment against high levels of IL-6 (Morozink, Friedman, Coe, & Ryff, 2010; see Boylan, Coe, & Ryff, Chapter 30, this volume). These characteristics, such as sense of control and psychological well-being, might be similarly protective for people who experience early adversity. We explore such possibilities in more detail in the next section below.

Future Directions

The research reviewed in this chapter highlights what research has already shown promotes resilience in terms of adult health among people who experience adversity in early life. This chapter also shows that there are many potentially fruitful areas for future research. We conclude by outlining additional questions that could be explored. Given that MIDUS is a large national sample that tracks a range of psychosocial and biological measures over time, it is especially well suited to studying questions about how early life adversity shapes the outcomes of diverse groups of people over time. Potential future topics for research include (a) investigation of additional traits that are likely to protect children experiencing adversity; (b) investigation of how buffers against the negative health effects of early life adversity might affect people differently, depending on their social or cultural backgrounds or on genetic factors; (c) investigation of how multiple psychosocial buffers of early life adversity might work together to protect health; and (d) tracking of individuals who have experienced early life adversity to assess health trajectories over time.

Other Protective Characteristics

As noted, there are additional psychological traits that might protect the health of people who have experienced adversity in early life but whose potential buffering effects have not yet been tested. Two examples are sense of control and psychological well-being. With respect to sense of control, while this construct has not been linked to better health among people experiencing early life adversity, a number of studies using data from MIDUS showed that sense of control can be protective for those who have lower SES in adulthood. For example, personal mastery, or the view that one has the ability to achieve one's goals, attenuates the relationship between income and self-rated health and acute symptoms (e.g., backaches, sleep difficulties; Lachman & Weaver, 1998). Perceived constraints, or the view that obstacles beyond one's control interfere with one's ability to achieve goals, play a similar role. Lack of perceived constraints buffers low-income individuals against the functional limitations (e.g., health-limiting ability to carry groceries or walk more than a mile) that are typically associated with adversity (Lachman & Weaver, 1998). Similarly, although people with lower levels of educational attainment are likely to die younger, research using MIDUS data showed that a strong sense of control buffers against this effect of lower levels of educational attainment on mortality (Turiano et al., 2014). Future research or other samples might test whether people who have experienced adversity in early life but who have a high sense of control in adulthood also have better health.

p. 56

Another individual psychological trait that is likely to protect the health of those who experienced adversity in early life is psychological well-being, which generally encompasses a variety of positive psychological states or the lack of negative psychological states. This can be measured in a variety of different ways, including lack of depression or negative affect, happiness or positive affect (hedonic well-being), or meaning in life and self-realization (eudaimonic well-being; Ryan & Deci, 2001; Ryff, 1989; Ryff & Keyes, 1995). While, again, no research that we are aware of has specifically explored whether psychological well-being protects against early life adversity in particular, there is evidence that it can buffer against adversity in adulthood. Using MIDUS, Morozink and colleagues found that lack of depression, positive affect, environmental mastery, positive relations with others, purpose in life, and self-acceptance all buffered individuals with lower levels of educational attainment against high levels of IL-6 (Morozink et al., 2010; see Boylan, Coe, & Ryff, Chapter 30, this volume). In fact, in many cases, the individuals with less education who had high levels of psychological well-being had levels of IL-6 that were comparable to those with more education. Future research with MIDUS or other samples could test whether people who have experienced adversity in early life but who have a high psychological well-being in adulthood also have better health.

Role of Demographic and Genetic Factors

Future research could also explore the ways that demographic and genetic factors influence the interaction between early life adversity and psychosocial buffers in determining adults' health. With respect to demographic characteristics, there are a number of differences that might emerge depending on gender, racial group membership, and the country in which one lives. Turning first to gender, previous research using MIDUS data has found that the relationships between early life adversity and adult health often differ between men and women (see Lee, Ryff, & Coe, Chapter 5, this volume). For example, among women, but not men, the link between childhood socioeconomic disadvantage and cardiovascular risk in adulthood is explained by having children at a younger age (Lee & Ryff, 2016). Thus, having higher levels of support in raising children might protect the health of women, in particular, when they have experienced adversity as children. Future research could investigate this idea in MIDUS by testing whether the relationship between childhood socioeconomic disadvantage and cardiovascular risk in adulthood is attenuated among women who receive assistance that might help them care or provide for their children (e.g., financial or unpaid assistance from parents or in-laws).

With respect to race, some research has compared the effect of similar experiences of childhood adversity on adult health outcomes in African Americans and whites (see Slopen et al., 2010, for an example with MIDUS). However, less research has compared whether the factors that buffer African Americans and other people of color are the same ones that do so for whites. For example, connections in the community can protect the health of those who experience adversity in childhood (Evans & Kutcher, 2011; Thorlindsson et al., 2012), and black churches are often an especially strong source of community for African Americans (McRae, Carey, & Anderson-Scott, 1998). Thus, religion or connection to a church community might be especially protective of the health of African Americans who experience early life adversity.

With respect to national culture, the Midlife in Japan (MIDJA) study, which includes many of the same psychosocial and biomarker measures as MIDUS in a sample of adults in the Tokyo metropolitan area, offers opportunities for research comparing the role of early life adversity in adult health across multiple cultures. Although the majority of the research conducted on the relationship between early life adversity and adult health has used Western samples, evidence suggests that similar relationships emerge in East Asian populations (see Fry, McCoy, & Swales, 2012, for a review). Additional research using MIDUS and MIDJA could explore whether the psychosocial factors that attenuate these relationships in the United States also do so in Japan.

Finally, comparisons using samples of twins and siblings in MIDUS might help to understand the role that genetic factors play in the health of people who experience adversity in early life. Previous research with these samples in MIDUS suggested that genetic effects on health may be greater in some circumstances (e.g., among people who have lower levels of education or income; Hamdi, Krueger, & South, 2015; Johnson & Krueger, 2005; but see also Krieger, Chen, Coull, & Selby, 2005; van den Berg, Doblhammer, & Christensen, 2009; van den Berg, Doblhammer-Reiter, & Christensen, 2011), but that genetics factors play less of a role in determining health outcomes among other groups of people (e.g., among people who have high levels of control; Johnson & Krueger, 2005). It is possible that early life adversity exacerbates genetic effects on health but that buffers such as strong social relationships reduce the role that genetics play in health, even among those who have grown up in adverse circumstances.

Joint Role of Multiple Psychosocial Buffers

Research on factors that protect the health of people who have experienced early life adversity has primarily studied one factor at a time. However, in reality, multiple buffers may occur simultaneously and work in concert with each other. The existing literature offers some evidence of such effects. For example, in a sample of healthy 13- to 16-year-olds, Chen and colleagues (2013) found that among lower SES children, having a role model was linked to lower levels of IL-6, and this relationship was partly explained by these children also engaging in shift-and-persist coping strategies. Thus, it seems that relationships with role models may help children to develop shift-and-persist coping strategies, which in turn protect their health, despite their lower SES. Research with MIDUS participants has not examined how multiple buffers might work simultaneously to protect the health of those who experienced early life adversity. However, there is some evidence from this sample that having a combination of multiple psychosocial resources (i.e., control beliefs, social support, and engagement in physical exercise) is related to better cognitive performance, and SES disparities in cognition are mitigated among those who have more of these resources (Agrigoroaei & Lachman, 2011).

There are at least four possible hypotheses about how multiple psychosocial factors might work together to protect the health of those who experienced adversity in early life. First, it is possible that one psychosocial buffer might mediate the relationship between another buffer and health among children who experience adversity. Chen and colleagues' (2013) findings that role models give rise to shift-and-persist strategies that then protect children's health is an example of such a relationship. Second, it is possible that psychosocial buffers have an additive relationship. For example, as in Agrigoroaei and Lachman's (2011) study, the combination of buffers such as sense of control, social support, and engaging in physical exercise might be protective. A third possibility is that buffers have multiplicative effects. For example, strong family relationships and high levels of social capital might each enhance the protective role that the other plays if children who are close to their parents receive an especially strong health benefit from their parents' ties to the community. A final possibility is that once a child who is experiencing adversity has one protective factor, having another makes a negligible difference. MIDUS would be a particularly useful sample in which to test such hypotheses due to its comprehensive set of such measures.

Longitudinal Assessments of People Who Experienced Early Life Adversity

A final area that could be pursued further involves tracking people over time to assess trajectories of early life adversity as individuals age. Most research that has studied psychosocial buffers against early life adversity has looked at physiological measures, such as inflammation, metabolic syndrome, and allostatic load, that indicate risk for chronic diseases of aging down the line. Continuing to track these participants over time could allow researchers to investigate whether these individuals are later diagnosed with chronic diseases such as cardiovascular disease or diabetes or whether they have higher mortality rates, all outcomes for which markers such as inflammation, metabolic syndrome, and allostatic load indicate risk (Danesh et al., 2004; Ford, 2005; Karlamangla, Singer, & Seeman, 2006). Future research might track, for example, whether people who have experienced adversity in early life but who also have the psychosocial buffers reviewed in the present chapter, are less likely to be diagnosed with chronic diseases of aging or have lower rates of mortality. These clinical effects might, for example, begin to emerge as individuals transition to old age. With respect to MIDUS in particular, continuing to follow these participants as they age would allow researchers to test more hypotheses about the psychosocial buffers of early life adversity on outcomes such as clinical diagnoses and mortality. Such an approach would allow researchers to understand more about how persistent the effects of early life adversity are across time and whether they have long-term effects that reach to the end of an individual's life.

Conclusion

p. 58

In sum, although a large body of research shows that most people who experience adversity in early life have worse health as adults, some who experience such adversity are resilient. Because it assesses such a range of psychosocial factors, in addition to biological outcomes, MIDUS has shed light on what might protect these individuals' health. Specifically, research using MIDUS data suggested that social relationships both in and outside of the family, as well as individual psychological characteristics such as positive coping strategies and psychological well-being, may protect the health of children experiencing adversity. Future research can help not only deepen our understanding of what helps children to live healthy lives despite adversity but also be used to help promote better health for all.

References

Agrigoroaei, S., & Lachman, M. E. (2011). Cognitive functioning in midlife and old age: Combined effects of psychosocial and behavioral factors. *Journals of Gerontology: Psychological Sciences*, 66B, i130–i140. doi:10.1093/geronb/gbr017

[Google Scholar](#) [WorldCat](#)

An, J. S., & Cooney, T. M. (2006). Psychological well-being in mid to late life: The role of generativity development and parent-child relationships across the lifespan. *International Journal of Behavioral Development*, 30, 410–421.

doi:10.1177/0165025406071489

[Google Scholar](#) [WorldCat](#)

Bonanno, G. A. (2004). Loss, trauma, and human resilience: Have we underestimated the human capacity to thrive after extremely aversive events? *American Psychologist*, 59, 20–208. doi:10.1037/0003-066X.59.1.20

[Google Scholar](#) [WorldCat](#)

Boynton-Jarrett, R., Ryan, L. M., Berkman, L. F., & Wright, R. J. (2008). Cumulative violence exposure and self-rated health: Longitudinal study of adolescents in the United States. *Pediatrics*, 122, 961–970. doi:10.1542/peds.2007-3063

[Google Scholar](#) [WorldCat](#)

Brady, S. S., & Matthews, K. A. (2002). The influence of socioeconomic status and ethnicity on adolescents' exposure to stressful life events. *Journal of Pediatric Psychology*, 27, 575–583. doi:10.1093/jpepsy/27.7.575

[Google Scholar](#) [WorldCat](#)

Braveman, P., & Barclay, C. (2009). Health disparities beginning in childhood: A life-course perspective. *Pediatrics*, 124, S163–S175. doi:10.1542/peds.2009-1100D

[Google Scholar](#) [WorldCat](#)

Brody, G. H., Lei, M. K., Chen, E., & Miller, G. E. (2014). Neighborhood poverty and allostatic load in African American youth. *Pediatrics*, 134, e1362–e1368. doi:10.1542/peds.2014-1395

[Google Scholar](#) [WorldCat](#)

Buka, S. L., Stichick, T. L., Birdthistle, I., & Earls, F. J. (2001). Youth exposure to violence: Prevalence, risks, and consequences. *American Journal of Orthopsychiatry*, 71, 298–310. doi:10.1037/0002-9432.71.3.298

[Google Scholar](#) [WorldCat](#)

Bures, R. M. (2003). Childhood residential stability and health at midlife. *American Journal of Public Health*, 93, 1144–1148. doi:10.2105/AJPH.93.7.1144

[Google Scholar](#) [WorldCat](#)

Carroll, J. E., Gruenewald, T. L., Taylor, S. E., Janicki-Deverts, D., Matthews, K. A., & Seeman, T. E. (2013). Childhood abuse, parental warmth, and adult multisystem biological risk in the Coronary Artery Risk Development in Young Adults study.

Proceedings of the National Academy of Sciences of the United States of America, 110, 17149–17153.

doi:10.1073/pnas.1315458110

[Google Scholar](#) [WorldCat](#)

Cassidy, J., & Shaver, P. R. (Eds.). (2008). *Human attachment: Theory, research, and clinical applications*. New York: Guilford Press.

[Google Scholar](#) [Google Preview](#) [WorldCat](#) [COPAC](#)

Chen, E., Lee, W. K., Cavey, L., & Ho, A. (2013). Role models and the psychological characteristics that buffer low-socioeconomic-status youth from cardiovascular risk. *Child Development*, 84, 1241–1252. doi:10.1111/cdev.12037

[Google Scholar](#) [WorldCat](#)

Chen, E., McLean, K. C., & Miller, G. E. (2015). Shift-and-persist strategies: Associations with socioeconomic status and the regulation of inflammation among adolescents and their parents. *Psychosomatic Medicine*, 77, 371–382.

[Google Scholar](#) [WorldCat](#)

Chen, E., & Miller, G. E. (2012). "Shift-and-persist" strategies: Why low socioeconomic status isn't always bad for health. *Perspectives on Psychological Science*, 7, 135–158. doi:10.1177/1745691612436694

[Google Scholar](#) [WorldCat](#)

Chen, E., Miller, G. E., Kober, M. S., & Cole, S. W. (2011). Maternal warmth buffers the effects of low early-life socioeconomic status on pro-inflammatory signaling in adulthood. *Molecular Psychiatry*, 16, 729–737. doi:10.1038/mp.2010.53

[Google Scholar](#) [WorldCat](#)

Chen, E., Miller, G. E., Lachman, M. E., Gruenewald, T. L., & Seeman, T. E. (2012). Protective factors for adults from low childhood socioeconomic circumstances: The benefits of shift-and-persist for allostatic load. *Psychosomatic Medicine*, 74, 178–186. doi:10.1097/PSY.0b013e31824206fd

[Google Scholar](#) [WorldCat](#)

Chen, E., Strunk, R. C., Trethewey, A., Schreier, H. M. C., Maharaj, N., & Miller, G. E. (2011). Resilience in low-socioeconomic-status children with asthma: Adaptations to stress. *Journal of Allergy and Clinical Immunology*, 128, 970–976. doi:10.1016/j.jaci.2011.06.040

[Google Scholar](#) [WorldCat](#)

Chen, E., Turiano, N. A., Mroczek, D. K., & Miller, G. E. (2016). Association of reports of childhood abuse and all-cause mortality rates in women. *JAMA Psychiatry*, 73, 920. doi:10.1001/jamapsychiatry.2016.1786

[Google Scholar](#) [WorldCat](#)

Cohen, S. (2004). Social relationships and health. *American Psychologist*, 59, 676–684. doi:10.1037/0003-066X.59.8.676

[Google Scholar](#) [WorldCat](#)

Cornier, M. A., Dabelea, D., Hernandez, T. L., Lindstrom, R. C., Steig, A. J., Stob, N. R., . . . Eckel, R. H. (2008). The metabolic syndrome. *Endocrine Reviews*, 29(7), 777–822. doi:10.1210/er.2008-0024

[Google Scholar](#) [WorldCat](#)

Danese, A., & McEwen, B. S. (2012). Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiology and Behavior*, 106, 29–39. doi:10.1016/j.physbeh.2011.08.019

[Google Scholar](#) [WorldCat](#)

Danesh, J., Wheeler, J. G., Hirschfield, G. M., Eda, S., Eiriksdottir, G., Rumley, A., Lowe, G. D. O., Pepys, M. B., & Gudnason, V. (2004). C-Reactive Protein and Other Circulating Markers of Inflammation in the Prediction of Coronary Heart Disease. *New England Journal of Medicine*, 350, 1387–1397. doi:10.1056/NEJMoa032804

[Google Scholar](#) [WorldCat](#)

Dong, M., Giles, W. H., Felitti, V. J., Dube, S. R., Williams, J. E., Chapman, D. P., & Anda, R. F. (2004). Insights into causal pathways for ischemic heart disease: Adverse Childhood Experiences Study. *Circulation*, 110, 1761–1766.

doi:10.1161/01.CIR.0000143074.54995.7F

[Google Scholar](#) [WorldCat](#)

Dunkel Schetter, C., & Dolbier, C. (2011). Resilience in the context of chronic stress and health in adults. *Social and Personality and Psychology Compass*, 5, 634–654. doi:10.1111/j.1751-9004.2011.00379.x

[Google Scholar](#) [WorldCat](#)

Epel, E., Jimenez, S., Brownell, K., Stroud, L., Stoney, C., & Niaura, R. (2004). Are stress eaters at risk for the metabolic syndrome? *Annals of the New York Academy of Sciences*, 1032, 208–210. doi:10.1196/annals.1314.022

[Google Scholar](#) [WorldCat](#)

Evans, G. W., Kim, P., Ting, A. H., Tessler, H. B., & Shannis, D. (2007). Cumulative risk, maternal responsiveness, and allostatic load among young adolescents. *Developmental Psychology*, 43, 341–351. doi:10.1037/0012-1649.43.2.341

Evans, G. W., & Kutcher, R. (2011). Loosening the link between childhood poverty and adolescent smoking and obesity: The protective effects of social capital. *Psychological Science, 22*, 3–7. doi:10.1177/0956797610390387

[Google Scholar](#) [WorldCat](#)

- p. 59 Fisher, P. A., Gunnar, M. R., Chamberlain, P., & Reid, J. B. (2000). Preventive intervention for maltreated preschool children: Impact on children's behavior, neuroendocrine activity, and foster parent functioning. *Journal of the American Academy of Child & Adolescent Psychiatry, 39*, 1356–1364. doi:10.1097/00004583-200011000-00009

[Google Scholar](#) [WorldCat](#)

Ford, E. S. (2005). Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: A summary of the evidence. *Diabetes Care, 28*, 1769–1778. doi:10.2337/diacare.28.7.1769

[Google Scholar](#) [WorldCat](#)

Friedman, E. M., Hayney, M., Love, G. D., Singer, B. H., & Ryff, C. D. (2007). Plasma interleukin-6 and soluble IL-6 receptors are associated with psychological well-being in aging women. *Health Psychology, 26*, 305–313. doi:10.1037/0278-6133.26.3.305

[Google Scholar](#) [WorldCat](#)

Friedman, E. M., Karlamangla, A. S., Gruenewald, T., Koretz, B., & Seeman, T. E. (2015). Early life adversity and adult biological risk profiles. *Psychosomatic Medicine, 77*, 176–185. doi:10.1097/PSY.0000000000000147

[Google Scholar](#) [WorldCat](#)

Friedman, E. M., Montez, J. K., Sheehan, C. M., Guenewald, T. L., & Seeman, T. E. (2015). Childhood adversities and adult cardiometabolic health: Does the quantity, timing, and type of adversity matter? *Journal of Aging and Health, 27*, 1311–1338.

doi:10.1177/0898264315580122

[Google Scholar](#) [WorldCat](#)

Fry, D., McCoy, A., & Swales, D. (2012). The consequences of maltreatment on children's lives: A systematic review of data from the East Asian and Pacific Region. *Trauma, Violence, and Abuse, 13*, 209–233. doi:10.1177/1524838012455873

[Google Scholar](#) [WorldCat](#)

Galobardes, B., Lynch, J. W., & Davey Smith, G. (2004). Childhood socioeconomic circumstances and cause-specific mortality in adulthood: Systematic review and interpretation. *Epidemiologic Reviews, 26*, 7–21. doi:10.1093/epirev/mxh008

[Google Scholar](#) [WorldCat](#)

Goodwin, R. D., & Wamboldt, F. S. (2012). Childhood physical abuse and respiratory disease in the community: The role of mental health and cigarette smoking. *Nicotine & Tobacco Research, 14*, 91–97. doi:10.1093/ntr/ntr126

[Google Scholar](#) [WorldCat](#)

Greenfield, E. A., & Marks, N. F. (2009a). Profiles of physical and psychological violence in childhood as a risk factor for poorer adult health: Evidence from the 1995–2005 National Survey of Midlife in the United States. *Journal of Aging and Health, 21*, 943–966. doi:10.1177/0898264309343905

[Google Scholar](#) [WorldCat](#)

Greenfield, E. A., & Marks, N. F. (2009b). Violence from parents in childhood and obesity in adulthood: Using food in response to stress as a mediator of risk. *Social Science & Medicine, 68*, 791–798. doi:10.1016/j.socscimed.2008.12.004

[Google Scholar](#) [WorldCat](#)

Greenfield, E. A., & Marks, N. F. (2010a). Identifying experiences of physical and psychological violence in childhood that jeopardize mental health in adulthood. *Child Abuse & Neglect, 34*, 161–171. doi:10.1016/j.chiabu.2009.08.012

[Google Scholar](#) [WorldCat](#)

Greenfield, E. A., & Marks, N. F. (2010b). Sense of community as a protective factor against long-term psychological effects of childhood violence. *Social Service Review, 84*, 129–147. doi:10.1086/652786

[Google Scholar](#) [WorldCat](#)

Gruenewald, T. L., Karlamangla, A. S., Hu, P., Stein-Merkin, S., Crandall, C., Koretz, B., & Seeman, T. E. (2012). History of socioeconomic disadvantage and allostatic load in later life. *Social Science & Medicine*, *74*, 75–83. doi:10.1016/j.socscimed.2011.09.037

[Google Scholar](#) [WorldCat](#)

Hamdi, N. R., Krueger, R. F., & South, S. C. (2015). Socioeconomic status moderates genetic and environmental effects on the amount of alcohol use. *Alcoholism: Clinical and Experimental Research*, *39*, 603–610. doi:10.1111/acer.12673

[Google Scholar](#) [WorldCat](#)

Hostinar, C. E., Lachman, M. E., Mroczek, D. K., Seeman, T. E., & Miller, G. E. (2015). Additive contributions of childhood adversity and recent stressors to inflammation at midlife: Findings from the MIDUS study. *Developmental Psychology*, *51*, 1630–1644. doi:10.1037/dev0000049

[Google Scholar](#) [WorldCat](#)

Hostinar, C. E., Ross, K. M., Chen, E., & Miller, G. E. (2017, April). *Social trust buffers against metabolic risk associated with childhood socioeconomic disadvantage*. Paper presented at the biannual meeting of the Society for Research in Child Development, Austin, TX.

Johnson, W., & Krueger, R. F. (2005). Higher perceived life control decreases genetic variance in physical health: Evidence from a national twin study. *Journal of Personality & Social Psychology*, *88*, 165–173. doi:10.1037/0022-3514.88.1.165

[Google Scholar](#) [WorldCat](#)

Juster, R. P., McEwen, B. S., & Lupien, S. J. (2010). Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience & Biobehavioral Reviews*, *35*, 2–16. doi:10.1016/j.neubiorev.2009.10.002

[Google Scholar](#) [WorldCat](#)

Kallem, S., Carroll-Scott, A., Rosenthal, L., Chen, E., Peters, S. M., McCaslin, C., & Ickovics, J. R. (2013). Shift-and-persist: A protective factor for elevated BMI among low-socioeconomic-status children. *Obesity*, *21*, 1759–1763. doi:10.1002/oby.20195

[Google Scholar](#) [WorldCat](#)

Karlamangla, A. S., Singer, B. H., & Seeman, T. E. (2006). Reduction in allostatic load in older adults is associated with lower all-cause mortality risk: MacArthur studies of successful aging. *Psychosomatic Medicine*, *68*, 500–507. doi:10.1097/01.psy.0000221270.93985.82

[Google Scholar](#) [WorldCat](#)

Keinan-Boker, L., Vin-Raviv, N., Liphshitz, I., Linn, S., & Barchana, M. (2009). Cancer incidence in Israeli Jewish survivors of World War II. *Journal of the National Cancer Institute*, *101*, 1489–1500. doi:10.1093/jnci/djp327

[Google Scholar](#) [WorldCat](#)

Kelly-Irving, M., Lepage, B., Dedieu, D., Bartley, M., Blane, D., Grosclaude, P., . . . Delpierre, C. (2013). Adverse childhood experiences and premature all-cause mortality. *European Journal of Epidemiology*, *28*, 721–734. doi:10.1007/s10654-013-9832-9

[Google Scholar](#) [WorldCat](#)

Krieger, N., Chen, J. T., Coull, B. A., & Selby, J. V. (2005). Lifetime socioeconomic position and twins' health: An analysis of 308 pairs of United States women twins. *PLoS Medicine*, *2*, e235. doi:10.1371/journal.pmed.0020235

[Google Scholar](#) [WorldCat](#)

Kubzansky, L. D., & Kawachi, I. (2000). Going to the heart of the matter: Do negative emotions cause coronary heart disease. *Journal of Psychosomatic Research*, *48*, 323–337. doi:10.1016/S0022-3999(99)00091-4

[Google Scholar](#) [WorldCat](#)

Laaksonen, M., Rahkonen, O., Martikainen, P., & Lahelma, E. (2005). Socioeconomic position and self-rated health: The contribution of childhood socioeconomic circumstances, adult socioeconomic status, and material resources. *American Journal*

of *Public Health*, 95, 1403–1409. doi:10.2105/AJPH.2004.047969

[Google Scholar](#) [WorldCat](#)

Lachman, M. E., & Weaver, S. L. (1998). The sense of control as a moderator of social class differences in health and well-being. *Journal of Personality and Social Psychology*, 74, 763–773. doi:10.1037/0022-3514.74.3.763

[Google Scholar](#) [WorldCat](#)

Lakka, H. M., Laaksonen, D. E., Lakka, T. A., Niskanen, L. K., Kumpusalo, E., Tuomilehto, J., & Kalonen, J. T. (2002). The metabolic syndrome and total and cardiovascular disease mortality in middle-aged men. *JAMA*, 288, 2709. doi:10.1001/jama.288.21.2709

[Google Scholar](#) [WorldCat](#)

Lee, C., & Ryff, C. D. (2016). Early parenthood as a link between childhood disadvantage and adult heart problems: A gender-based approach. *Social Science and Medicine*, 171, 58–66. doi:10.1016/j.socscimed.2016.10.028

[Google Scholar](#) [WorldCat](#)

p. 60 Lee, C., Tsenkova, V., & Carr, D. (2014). Childhood trauma and metabolic syndrome in men and women. *Social Science & Medicine*, 105, 122–130. doi:10.1016/j.socscimed.2014.01.017

[Google Scholar](#) [WorldCat](#)

Luthar, S. S. (2006). Resilience in development: A synthesis of research across five decades. In D. Cicchetti & D. J. Cohen (Eds.), *Developmental psychopathology: Vol. 3. Risk, disorder, and adaptation* (pp. 739–795). New York: Wiley.

[Google Scholar](#) [Google Preview](#) [WorldCat](#) [COPAC](#)

Maier, E. H., & Lachman, M. E. (2000). Consequences of early parental loss and separation for health and well-being in midlife. *International Journal of Behavioral Development*, 24, 183–189. doi:10.1080/016502500383304

[Google Scholar](#) [WorldCat](#)

Mallers, M. H., Charles, S. T., Neupert, S. D., & Almeida, D. M. (2010). Perceptions of childhood relationships with mother and father: Daily emotional and stressor experiences in adulthood. *Developmental Psychology*, 46, 1651–1661. doi:10.1037/a0021020

[Google Scholar](#) [WorldCat](#)

Masten, A. S. (2001). Ordinary magic: Resilience processes in development. *American Psychologist*, 56, 227–238.

doi:10.1037/0003-066X.56.3.227

[Google Scholar](#) [WorldCat](#)

McRae, M. B., Carey, P. M., & Anderson-Scott, R. (1998). Black churches as therapeutic systems: A group process perspective. *Health Education and Behavior*, 25, 778–789. doi:10.1177/109019819802500607

[Google Scholar](#) [WorldCat](#)

Miller, G. E., Brody, G. H., Yu, T., & Chen, E. (2014). A family-oriented psychosocial intervention reduces inflammation in low-SES African American youth. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 11287–11292.

doi:10.1073/pnas.1406578111

[Google Scholar](#) [WorldCat](#)

Miller, G. E., Chen, E., & Parker, K. J. (2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: Moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*, 137, 959–997. doi:10.1037/a0024768

[Google Scholar](#) [WorldCat](#)

Miller, G. E., Lachman, M. E., Chen, E., Gruenewald, T. L., Karlamangla, A. S., & Seeman, T. E. (2011). Pathways to resilience: Maternal nurturance as a buffer against the effects of childhood poverty on metabolic syndrome at midlife. *Psychological Science*, 22, 1591–1599. doi:10.1177/0956797611419170

doi:10.1177/0956797611419170

[Google Scholar](#) [WorldCat](#)

Moroziuk, J. A., Friedman, E. A., Coe, C. L., & Ryff, C. D. (2010). Socioeconomic and psychosocial predictors of interleukin-6 in the MIDUS national sample. *Health Psychology*, 29, 626–635. doi:10.1037/a0021360

[Google Scholar](#) [WorldCat](#)

Morton, P. M., Schafer, M. H., & Ferraro, K. F. (2012). Does childhood misfortune increase cancer risk in adulthood? *Journal of Aging and Health*, *24*, 948–984. doi:10.1177/0898264312449184

[Google Scholar](#) [WorldCat](#)

Oliver, G., Wardle, J., & Gibson, E. L. (2000). Stress and food choice: A laboratory study. *Psychosomatic Medicine*, *62*, 853–865. doi:10.1097/00006842-200011000-00016

[Google Scholar](#) [WorldCat](#)

Pressman, S. D., & Cohen, S. (2005). Does positive affect influence health? *Psychological Bulletin*, *131*, 925–971. doi:10.1037/0033-2909.131.6.925

[Google Scholar](#) [WorldCat](#)

Repetti, R. L., Robles, T. F., & Reynolds, B. (2011). Allostatic processes in the family. *Development and Psychopathology*, *23*, 921–938. doi:10.1017/S095457941100040X

[Google Scholar](#) [WorldCat](#)

Repetti, R. L., Taylor, S. E., & Seeman, T. E. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, *128*, 330–366. doi:10.1037/0033-2909.128.2.330

[Google Scholar](#) [WorldCat](#)

Ridker, P. M., Hennekens, C. H., Buring, J. E., & Rifai, N. (2000). C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *New England Journal of Medicine*, *342*, 836–843.

doi:10.1056/NEJM200003233421202

[Google Scholar](#) [WorldCat](#)

Ridker, P. M., Rifai, N., Stampfer, M. J., & Hennekens, C. H. (2000). Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation*, *101*, 1767–1772. doi:10.1161/01.CIR.101.15.1767

[Google Scholar](#) [WorldCat](#)

Rothrauff, T. C., Cooney, T. M., & An, J. S. (2009). Remembered parenting styles and adjustment in middle and late adulthood. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, *64B*, 137–146. doi:10.1093/geronb/gbn008

[Google Scholar](#) [WorldCat](#)

Ryan, R. M., & Deci, E. L. (2001). On happiness and human potentials: A review of research on hedonic and eudaimonic well-being. *Annual Review of Psychology*, *52*, 141–166. doi:10.1146/annurev.psych.52.1.141

[Google Scholar](#) [WorldCat](#)

Ryff, C. D. (1989). Happiness is everything, or is it? Explorations on the meaning of psychological well-being. *Journal of Personality and Social Psychology*, *57*, 1069–1081. doi:10.1037/0022-3514.57.6.1069

[Google Scholar](#) [WorldCat](#)

Ryff, C. D., & Keyes, C. L. M. (1995). The structure of psychological well-being revisited. *Journal of Personality and Social Psychology*, *69*, 719–727. doi:10.1037/0022-3514.69.4.719

[Google Scholar](#) [WorldCat](#)

Ryff, C. D., Singer, B. H., & Love, G. D. (2004). Positive health: Connecting well-being with biology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *359*, 1383–1394. doi:10.1098/rstb.2004.1521

[Google Scholar](#) [WorldCat](#)

Schafer, M. H., Morton, P. M., & Ferraro, K. F. (2014). Child maltreatment and adult health in a national sample: Heterogeneous relational contexts, divergent effects? *Child Abuse & Neglect*, *38*, 395–406. doi:10.1016/j.chiabu.2013.08.003

[Google Scholar](#) [WorldCat](#)

Schofield, T. J., Conger, R. D., Gonzales, J. E., & Merrick, M. T. (2016). Harsh parenting, physical health, and the protective role of positive parent-adolescent relationships. *Social Science & Medicine*, *157*, 18–26. doi:10.1016/j.socscimed.2016.03.027

[Google Scholar](#) [WorldCat](#)

Schrepf, A., Markon, K., & Lutgendorf, S. K. (2014). From childhood trauma to elevated C-reactive protein in adulthood: The role of anxiety and emotional eating. *Psychosomatic Medicine*, *76*, 327–336. doi:10.1097/PSY.0000000000000072

[Google Scholar](#) [WorldCat](#)

Seeman, T., Epel, E., Gruenewald, T., Karlamangla, A., & McEwen, B. S. (2010). Socio-economic differentials in peripheral biology: Cumulative allostatic load: SES peripheral biology. *Annals of the New York Academy of Sciences*, *1186*, 223–239.

doi:10.1111/j.1749-6632.2009.05341.x

[Google Scholar](#) [WorldCat](#)

Seeman, T. E., McEwen, B. S., Rowe, J. W., & Singer, B. H. (2001). Allostatic load as a marker of cumulative biological risk: MacArthur studies of successful aging. *Proceedings of the National Academy of Sciences of the United States of America*, *98*, 4770–4775. doi:10.1073/pnas.081072698

[Google Scholar](#) [WorldCat](#)

Shonkoff, J. P., Boyce, W. T., & McEwen, B. S. (2009). Neuroscience, molecular biology, and the childhood roots of health disparities: Building a new framework for health promotion and disease prevention. *JAMA*, *301*, 2252.

doi:10.1001/jama.2009.754

[Google Scholar](#) [WorldCat](#)

Shonkoff, J. P., & Garner, A. S. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, *129*, e232–e246.

doi:10.1542/peds.2011-2663

[Google Scholar](#) [WorldCat](#)

Slopen, N., Lewis, T. T., Gruenewald, T. L., Mujahid, M. S., Ryff, C. D., Albert, M. A., & Williams, D. R. (2010). Early life adversity and inflammation in African Americans and whites in the Midlife in the United States survey. *Psychosomatic Medicine*, *72*, 694–701. doi:10.1097/PSY.0b013e3181e9c16f

Taha, F., Galea, S., Hien, D., & Goodwin, R. D. (2014). Childhood maltreatment and the persistence of smoking: A longitudinal study among adults in the US. *Child Abuse & Neglect*, *38*, 1995–2006. doi:10.1016/j.chiabu.2014.10.022

[Google Scholar](#) [WorldCat](#)

Taylor, S. E. (2010). Mechanisms linking early life stress to adult health outcomes. *Proceedings of the National Academy of Sciences of the United States of America*, *107*, 8507–8512. doi:10.1073/pnas.1003890107

[Google Scholar](#) [WorldCat](#)

Thorlindsson, T., Valdimarsdottir, M., & Hrafn Jonsson, S. (2012). Community social structure, social capital and adolescent smoking: A multi-level analysis. *Health & Place*, *18*, 796–804. doi:10.1016/j.healthplace.2012.03.013

[Google Scholar](#) [WorldCat](#)

Tsenkova, V., Boylan, J. M., & Ryff, C. (2013). Stress eating and health: Findings from MIDUS, a national study of US adults. *Appetite*, *69*, 151–155. doi:10.1016/j.appet.2013.05.020

[Google Scholar](#) [WorldCat](#)

Tsenkova, V., Pudrovska, T., & Karlamangla, A. (2014). Childhood socioeconomic disadvantage and prediabetes and diabetes in later life: a study of biopsychosocial pathways. *Psychosomatic Medicine*, *76*, 622–628. doi:10.1097/PSY.0000000000000106

[Google Scholar](#) [WorldCat](#)

Turiano, N. A., Chapman, B. P., Agrigoroaei, S., Infurna, F. J., & Lachman, M. (2014). Perceived control reduces mortality risk at low, not high, education levels. *Health Psychology*, *33*, 883–890. doi:10.10w37/hea0000022

[Google Scholar](#) [WorldCat](#)

Uchino, B. N., Cacioppo, J. T., & Kiecolt-Glaser, J. K. (1996). The relationship between social support and physiological processes: A review with emphasis on underlying mechanisms and implications for health. *Psychological Bulletin*, *119*, 488–531. doi:10.1037/0033-2909.119.3.488

[Google Scholar](#) [WorldCat](#)

Uphold-Carrier, H., & Utz, R. (2012). Parental divorce among young and adult children: A long-term quantitative analysis of mental health and family solidarity. *Journal of Divorce & Remarriage*, *53*, 247–266. doi:10.1080/10502556.2012.663272

[Google Scholar](#) [WorldCat](#)

van den Berg, G. J., Doblhammer, G., & Christensen, K. (2009). Exogenous determinants of early-life conditions, and mortality later in life. *Social Science and Medicine*, *68*, 1591–1598. doi:10.1016/j.socscimed.2009.02.007

[Google Scholar](#) [WorldCat](#)

van den Berg, G. J., Doblhammer-Reiter, G., & Christensen, K. (2011). Being born under adverse economic conditions leads to a higher cardiovascular mortality rate later in life: Evidence based on individuals born at different stages of the business cycle. *Demography*, *48*, 507–530. doi:10.1007/s13524-011-0021-8

[Google Scholar](#) [WorldCat](#)

Wegman, H. L., & Stetler, C. (2009). A meta-analytic review of the effects of childhood abuse on medical outcomes in adulthood: *Psychosomatic Medicine*, *71*, 805–812. doi:10.1097/PSY.0b013e3181bb2b46

p. 62

[Google Scholar](#) [WorldCat](#)